

BMJ Open

The Canadian Survey of Health, Lifestyle and Aging with Multiple Sclerosis; Methodology and Initial Results

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2014-005718
Article Type:	Research
Date Submitted by the Author:	20-May-2014
Complete List of Authors:	Ploughman, Michelle; Memorial University of Newfoundland, Faculty of Medicine Beaulieu, Serge; Eastern Health Authority, Harris, Chelsea; Memorial University, Faculty of Medicine Hogan, Stephen; Memorial University, Faculty of Medicine Manning, Olivia; Memorial University, Faculty of Medicine Alderdice, Penelope; Memorial University, Faculty of Medicine Fisk, John; Dalhousie University, Sadovnick, A; University of British Columbia, Department of Medical Genetics and Division of Neurology O'Connor, Paul; St. Michaels Regional Hospital, Department of Neurology Morrow, Sarah; London Health Science Centre, Metz, Luanne; University of Alberta, Department of Neurology Smyth, Penelope; University of Calgary, Clinical Neurosciences Mayo, Nancy; McGill University, Medicine Marrie, Ruth; University of Manitoba, Departments of Internal Medicine and Community Health Sciences Knox, Katherine; University of Saskatchewan, Department of Physical Medicine and Rehabilitation Stefanelli, Mark; Memorial University, Faculty of Medicine Godwin, Marshall; Memorial University, Primary Health Care Research Unit
Primary Subject Heading:	Neurology
Secondary Subject Heading:	Research methods
Keywords:	Multiple sclerosis < NEUROLOGY, REHABILITATION MEDICINE, NEUROLOGY

SCHOLARONE™
Manuscripts

The Canadian Survey of Health, Lifestyle and Aging with Multiple Sclerosis; Methodology and Initial Results

Michelle Ploughman PT, PhD*, Serge Beaulieu**, Chelsea Harris BKin MPH*, Stephen Hogan BKin*, Olivia J. Manning*, Penelope W. Alderdice PhD*, John D. Fisk PhD[§], A. Dossa Sadovnick PhD[†], Paul O'Connor, MD, FRCPC[‡], Sarah A. Morrow, MD[†], FRCPC, Luanne M. Metz MD, FRCPC[#], Penelope Smyth MD, FRCPC[¶], Nancy Mayo, PhD^{§§}, Ruth Ann Marrie, MD, PhD^{††}, Katherine B. Knox, MD, FRCPC^{‡‡}, Mark Stefanelli MD FRCPC^{¶¶}, Dr. Marshall Godwin, MD^{##}

*Recovery and Performance Laboratory, Rehabilitation Research Unit, Faculty of Medicine, Memorial University, St. John's NL Canada

**Eastern Health Authority, St. John's NL Canada

§Dalhousie University, Halifax NS, Canada

†Department of Medical Genetics and Division of Neurology, Faculty of Medicine University of British Columbia, Vancouver BC, Canada

‡Department of Neurology, St. Michaels Regional Hospital, Toronto ON, Canada

†London Health Science Centre, London ON, Canada

¶Department of Neurology, University of Alberta, Edmonton AB, Canada

¶Clinical Neurosciences, University of Calgary, Calgary AB, Canada

§§Clinical Epidemiology, McGill University, Montreal QC, Canada

††Departments of Internal Medicine and Community Health Sciences, University of Manitoba, Winnipeg MB, Canada

‡‡Department of Physical Medicine and Rehabilitation, College of Medicine, University of Saskatchewan, Saskatoon SK, Canada

¶¶Neurology, Faculty of Medicine, Memorial University, St. John's NL Canada

##Primary Health Care Research Unit, Faculty of Medicine, Memorial University, St. John's NL, Canada

Corresponding Author/Research Site

Dr. Michelle Ploughman, Recovery and Performance Laboratory, Rehabilitation Research Unit, Faculty of Medicine, Memorial University, Rm 400, L.A. Miller Centre, 100 Forest Rd., St. John's NL Canada. Ph 709 777 2099; fax 709 777 8702; email mploughm@mun.ca

Key Words: aging; diet; exercise; health behaviours; multiple sclerosis; neurodegenerative disease; older persons; quality of life

Short Title: Health Lifestyle and Aging with Multiple Sclerosis

Tables: 3

Figures: 2

For peer review only

ABSTRACT

Objective

People with multiple sclerosis (MS) are living longer so strategies to enhance long term health and quality of life are garnering more interest. We aimed to create a profile of aging with MS in Canada focusing on health and lifestyle factors, disability, participation and quality of life in order to determine factors associated with healthy aging.

Design

National multi-centre postal survey

Setting

Recruitment from Canadian MS clinics, MS Society of Canada chapters, and newspaper advertisements

Participants

People aged 55 years or older with MS symptoms more than 20 years

Outcome Measures

Validated outcome measures and custom-designed questions examining MS disease characteristics, living situation, disability, co-morbid conditions, fatigue, health behaviours, mental health, social support, impact of MS, and others.

Results

Of the 921 surveys, 743 were returned (80.7% response rate). Participants (mean age 64.6 ±6.2 years) reported living with MS symptoms for an average of 32.9±9.5 years and 28.6% were either wheelchair users or bedridden. There was only 5.4% missing data and 709 respondents provided optional qualitative information. According to data derived from the 2012 Canadian Community Health Survey of Canadians over 55 years of age, older people with MS from this survey sample are about 8 times less likely to be employed full-time. Older people with MS were less likely to engage in regular physical activity (26.7%) compared to typical older Canadians (45.2%). However, they were more likely to abstain from alcohol and smoking.

Conclusion

Despite barriers to participation, we were able to recruit and gather detailed responses (with good data quality) from a large proportion of older Canadians with MS. The data suggests that this sample of older people with MS are less likely to be employed, are less active and more disabled than other older Canadians.

Strengths and limitations of this study

- This unique cohort provides new insights on aging with MS. With an average age about 65 years, our respondents are 10 to 25 years older than cohorts examined in previous studies of older people with MS.
- Our recruitment methods and a very high survey return rate allowed us to obtain data from about 3% of our target population.
- With only about 10% of our sample being employed in any capacity, our results point to a need to understand retirement and employment in aging with MS.
- The cross sectional design limits our ability to assess change and the effects of variables on individual outcomes over time.
- The low sampling of persons in nursing homes and assisted living situations highlights the need for specialized recruitment strategies to study this sub-set of people living with MS.

This research was supported by a Canadian Institutes of Health Research (CIHR) Post-doctoral Fellowship, a Newfoundland and Labrador Centre for Applied Health Research, Healthy Aging Research Program Project Grant, the Health Care Foundation (Eastern Health Authority) Research Project Grant, and the Physiotherapy Foundation of Canada Alberta Research Award.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Multiple Sclerosis (MS) is a highly unpredictable and heterogeneous autoimmune disease diagnosed between the ages of 18-40 years. Affecting women more than men, the cause of MS is unknown, however research suggests that a combination of environmental factors and genetic predisposition may be the trigger for disease onset [1]. The number of people with MS over the age of 55 is increasing, possibly due to improved longevity and more effective treatments [2]. On the other hand, several authors have shown that smoking [3] and cardiovascular disease [4] accelerate disease progression. People with MS are interested in strategies that minimize disability and enhance role participation and quality of life [5]. The uncertainty surrounding MS progression along with aging MS patients warrants the need for investigation into potential disease modifying health behaviours. Older individuals who have lived with MS for many years may provide insight into factors that contribute to living a long and healthy life with MS [6].

Older people with MS are often excluded from MS research. Studies involving clinical drug trials [7] and exercise interventions [8], in particular, target patients with mild relapsing-remitting disease. Furthermore, because they may not require intensive medical oversight associated with taking disease-modifying drugs, older patients may not attend MS clinics as often as those recently diagnosed. The older MS cohort are also more likely to be disabled and living in an institutional setting, making recruitment challenging. In addition, disability-related impairments such as those affecting vision, cognition and coordination complicate survey and patient-reported outcome administration [9].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In order to create a profile of aging with MS in Canada, we developed a national cross-sectional survey, based on previous qualitative research [5 6], using validated patient-reported outcome measures [9]. We focused on health and lifestyle factors, disability, participation and quality of life, targeting people 55 years of age and older with MS symptoms for 20 or more years. By partnering with major MS clinic registries and MS society chapters, we hypothesized that we would be able to reach a representative sample of about 5% of the ‘aging with MS in Canada’ cohort.

METHODS

Participants and Sampling: We targeted Canadians, aged 55 years or older, who had had self-reported MS symptoms for 20 or more years. Prior to recruitment, the study was approved by 11 health research ethics boards in eight Canadian provinces. Between May 2011 and December 2012, participants were recruited by telephone from lists of eligible MS clinic patients and by public advertising through the MS Society of Canada Research Internet Portal[10], local MS Society chapters and their newsletters, and newspaper advertisements. A confirmed diagnosis from a neurologist was available for participants recruited through the MS clinic databases; self-identified participants provided a diagnosis date and neurologist name to confirm their diagnosis. Rather than calculating a sample size, we used an exhaustive sampling technique to obtain a large and representative sample; for example repeatedly targeting regional newspapers, following-up at study sites, making formal and informal contacts with MS Society Chapter support groups.

After obtaining consent using a standard script, we mailed participants the questionnaire, with an information sheet, a toll-free telephone helpline number and email address and a prepaid

reply envelope. We requested that participants return the completed survey in 2 weeks; non-responders were telephoned for follow-up at 4 weeks. Participants who wished to complete the survey by telephone were encouraged to do so. Each questionnaire was coded to maintain anonymity.

Survey Design: The questionnaire consisted of validated patient-reported outcome tools and additional custom-designed questions chosen based on pilot research in a small sample of older people with MS [9]. Based on pilot work and participant feedback, a (paper) mail version of the survey using a larger 14-point font was divided into sections that grouped similar questions together. Respondents were instructed to complete the survey in these sections (chapters) taking a break in between. Since we previously observed confusion with questions involving temporal parameters, questions regarding activity involvement that followed the same timeline (e.g. weekly, monthly, or yearly) were placed in sequence. To further examine the acceptability of the survey and to minimize skipped or misunderstood items, we asked the first 14 participants (11 returns) to provide feedback on the content, format and response burden. This feedback was integrated into the final version of the questionnaire but the responses were not included in the final data.

In order to include people with MS in the province of Quebec, Canada, the entire survey including the cover letter was translated into french by one independent professional translator. We also obtained french versions of previously validated tools. Four native french speakers involved in MS research made corrections to the initial translation in sequence. The corrections were then consolidated by a fifth french speaking MS researcher. The advertisement and

consent scripts were translated by a native french speaking MS researcher and reviewed and corrected by a second.

Survey Content: The final survey consisted of five chapters (**Table 1**) ending with open response questions for participants to provide additional comments. Where appropriate, we obtained permission from the developers to use previously validated tools.

We used the Multiple Sclerosis Impact Scale (MSIS-29), the Barthel Index and the Co-morbidity Questionnaire to assess health. The MSIS-29 consists of 29 questions: 20 address the physical impact and nine address the psychological impact of MS generating separate or combined scores as a percentage of 100. Higher scores indicate greater impact of MS [11]. The Barthel Index measures function and mobility in activities of daily living (ADL) in ten areas: feeding, bathing, grooming, dressing, bowel control, bladder control, toileting, transfers, ambulation and stair climbing [12]. Items are rated in terms of whether individuals can perform activities independently, with some assistance, or are completely dependent, with a total score ranging from 0 (complete assistance required) to 100 (independent). The Barthel Index has been tested in people with MS [13-15] and is reliable whether administered by telephone, in-person or by mail [16]. The list of co-morbidities used in this survey was adapted from Marrie and Horwitz [17 18] in consultation with Dr. Marrie (personal communication).

We determined the degree of participation in instrumental activities of daily living (iADL) using the Frenchay Activities Index (FAI) and individual components of lifestyle (exercise, diet, smoking, alcohol, and stress) using the Simple Lifestyle Indicator Questionnaire

(SLIQ). The FAI assesses the frequency of participation in domestic chores, social activities, work and leisure for a summed score from 0 – 45; higher score indicating more frequent participation [19]. The FAI has excellent test-retest reliability as a mailed questionnaire [20] and correlates well with face-to-face interview scores [21]. The FAI has been mainly used in stroke but has also been used in large MS population studies [22 23]. The SLIQ was developed to measure the individual components of lifestyle that affect cardiovascular disease[24]. Initial content and face validity were assessed by the developers among 136 people attending a family practice clinic[24].

We were also interested in mental health, coping and social support in aging with MS. The Hospital Anxiety and Depression Scale (HADS) is a self-assessment scale found to reliably screen for depression and anxiety in multiple sclerosis patients [25]. It is divided into an Anxiety subscale (HADS-A) and a Depression subscale (HADS-D) each containing seven intermingled items scored out of 21; higher score indicating greater symptoms [26]. Resilience, the ability to overcome adversity, was measured using the 25 item Resilience Scale. Responses to the test questions range from 1 (disagree) to 7 (agree) and the summed score can be further interpreted as, low resilience (<120), moderate resilience (125-145), and high resilience (>145) [27] and has been validated in older community-living individuals [27]. The Personal Resource Questionnaire (PRQ-2000) measures social support and consists of 15 statements with which the subject agrees or disagrees based on a seven-point Likert scale; higher scores indicating higher support (e.g. I belong to a group in which I feel important 1-Strongly Disagree to 7-Strongly Agree). PRQ-2000 demonstrates adequate content, predictive and construct validity [28] and mailed versions have been used in community-living women with chronic health conditions[29] and adults with MS[30].

Table 1: Survey Content

CHAPTER	DOMAINS	MEASUREMENT TOOLS
You and Your MS (5 pages)	Demographics	Postal code, gender, height, weight, age, finances and education
	MS Disease Characteristics	Symptom onset, time since diagnosis, and type of MS
	Environment	Living situation, accessibility and home modifications
Your Health (6 pages)	Health-Related Quality of Life	Visual analogue scale of perceived health status [9] Multiple Sclerosis Impact Scale-29 [11]
	Physical Health and Disability	Barthel Index [12] Co-Morbidity Questionnaire [17]
	Health Care Services	Use and satisfaction with health care services, medications and complementary/alternative health.
Your Activities (7 pages)	Participation and instrumental Activities of Daily Living	Frenchay Activities index [19]; Employment and impact of fatigue
	Lifestyle	Simple Lifestyle Indicator Questionnaire (diet, exercise, stress, smoking, alcohol) [24]
Your Thoughts and Feelings (8 pages)	Mental/ Cognitive Health	Hospital Anxiety and Depression Scale (HADS) Resilience Scale [25] Problems with concentration and memory
Your Social Life (4 pages)	Social Support	Personal Resource Questionnaire-2000 [30]

Canadian comparison data: Comparative data was collected from the Canadian Community Health Survey (CCHS) 2012 version; a national cross-sectional survey that gathers information about health, social and economic status and lifestyle behaviours of the population. Data was extracted from the CCHS database using the University of Toronto Data Library Service (DLS) [31]. A CCHS sample was selected based on age (>55 years). Data was collected from CCHS questions that matched question from our survey. Outputs (ie. highest level of education, employment status) were analyzed using Survey Documentation & Analysis (SDA) software [32]. Frequency tables were created for variables of interest.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Data Entry: We custom designed the database application to house project data and minimize error during data imputation. The application was constructed using Borland Delphi 2005 as the application interface (front end) and a freeware (<5 user) version of the Advantage database engine for the back end. To facilitate reporting and publication, an export routine was programmed into the database to perform the following functions: Generate a pipe delimited text file of the main database table, iterate through the application, and associated database lookup tables, and generate an SPSS syntax file with the applicable variable and value labels. The pipe delimited text file was imported into SPSS and an SPSS syntax file of the import was generated for recoding. An SPSS syntax file was generated for recoding data according to the needs outlined in the scoring tool and applied to the data and reused with each data dump. Every analysis performed was saved as a syntax file in order to facilitate replication, documentation and publication.

Data Cleaning and Imputation: Missing data was imputed using three techniques: (1) deductive imputation, (2) person mean substitution, and (3) ‘hot deck’ imputation [33]. The deductive technique was used to impute missing data when there was sufficient comments and answers to related questions. If less than half of the data was missing on a section, scale, or subscale, person mean substitution (PMS) was used. PMS takes the mean score of the particular participant on that section, scale, or subscale and uses the mean as the value for the missing data. If greater than half of the data was missing on a section, scale, or subscale, ‘hot deck’ imputation was used [33]. This involved taking a value from the "hot deck" or in other terms, finding another participant of the same gender, age (± 5 years), education level, and living in the same province. Hot deck

imputation was selected for this study because it is proven to be the most efficient method of data imputation as it is more robust in large samples when compared to other imputation methods [33]. The exact scores for each question from the substitute participant were imputed for the missing values for the participant of interest. Variables that could not be imputed and remained empty included, from **Table 1**, Demographics (e.g. weight, height, finances), MS Disease Characteristics, Environment, and Health Care Services.

Data Analysis: Data was analyzed using IBM SPSS software v20. Frequencies and descriptive statistics (mean, SD and data range) were used to determine subject characteristics.

RESULTS

Survey Response

Seven hundred and forty three participants returned the survey (**Figure 1**; 80.7% response rate). Survey response by province is illustrated in **Figure 2**. Of the respondents, 90.4% (n=672) were identified through provincial MS clinics while the remaining 71 self-identified by responding to newspaper ads and word of mouth. Forty-five people requested help to complete the survey, and 11 participants asked that the entire survey be completed by phone due to writing or vision difficulties or both. Additional content was provided by 709 participants in the open-ended questions at the end of the survey.

Only 563 of 104,763 cells contained missing data (5.37%). Almost every survey contained one or two random missing items; in some cases answers to these could be inferred using deductive reasoning ($67/104,763=0.06\%$). We observed no pattern of missing data other

than body weight (which we did not impute). The PMS technique was required for 170 cells (170/94,763=0.16%) and the ‘hot deck’ technique was required in 14 surveys (324 cells/104,763=0.31%). Cases of consecutive missing data seemed to be mainly due to pages adhering together resulting in questions being overlooked.

Insert Figure 1 about here

Sample Characteristics

Participants ranged in age from 55 to 88 years with a mean age of 64.6 years (± 6.18). They described experiencing MS symptoms for 20 to 73 years with a mean duration of 32.9 years (± 9.5). The respondents reported that the average time from onset of symptoms to diagnosis by a neurologist was 8.05 years (± 9.39) with a range from 0 to 47 years. The number of women outnumbered men 3.48:1 (**Table 2**).

Sixty percent reported at least one year of post-secondary education, and 90% were either retired or unemployed. About twenty-eight percent of the sample (n=212) reported that they were no longer able to walk and either used a wheelchair or were bedridden. The characteristics of the sample are fully described in **Table 2**.

Table 2: Sample Characteristics

Characteristic		N (%)
Gender	Males	166 (22.3)
	Females	577 (77.7)
Age	55-64	414 (55.7)
	65-74	274 (36.9)
	75-84	50 (6.7)
	85+	5 (0.7)
Years with MS Symptoms	20-29	324 (43.6)
	30-39	245 (33.0)
	40-49	135 (18.2)
	50-59	31 (4.2)
	60 or more	7 (0.9)
Initial MS Diagnosis	Relapsing-Remitting	386 (52.0)
	Primary Progressive	99 (13.3)
	Secondary Progressive	65 (8.7)
	Progressive-Relapsing	16 (2.2)
	Benign	43 (5.8)
	Unknown	126 (17.0)
Living situation	Living in a private home	715 (96.2)
	Living in an assisted living complex	16 (2.2)
	Living in long term care	11 (1.5)
	Home Care	146 (19.7)
	Equipment	200 (26.9)

	Home Modifications	73 (9.8)
Private Insurance (Health/Disability)	Yes	393 (52.9)
	No	350 (47.1)

Insert Figure 2 about here

Older people with MS compared to typical older Canadians

According to data derived from the 2012 Canadian Community Health Survey (CCHS; **Table 3**) of Canadians over 55 years of age, older people with MS from this survey sample have less post-secondary education and are about 8 times less likely to be employed full-time. About half of the survey sample have major mobility challenges (unable to walk outdoors, use a wheelchair or are bedridden) compared to only 9% in the CCHS sample. Not surprising, older people with MS were less likely to engage in regular physical activity (26.7%) compared to typical older Canadians (45.2%). However, they were more likely to abstain from alcohol and smoking.

Table 3: Comparison of Sample Characteristics to Others Aging in Canada

Characteristic		Study sample (%)	CCHS sample (%)
Education	Less than high school	74 (10.0)	1,241,946 (15.2)
	High School	222 (29.9)	1,083,618 (13.3)
	1-2 year post-secondary	189 (25.4)	4,088,252 (4.9)
	Complete Post-graduate	258 (34.7)	5,442,586 (66.6)
Employment	Retired	504 (67.8)	2,878,200 (46.3)

Status	Unemployed	166 (22.3)	302,942 (4.8)
	Employed Full-time	30 (4.0)	2,068,571 (33.3)
	Employed Part-time	19 (2.6)	698,125 (11.2)
	Employed Casual	24 (3.2)	266,419 (4.3)
Mobility	Walk independently with or without cane outdoors	351 (47.2)	8,004,425 (90.9)
	Walk with cane or help indoors	180 (24.2)	593,584 (6.7)
	Wheelchair	161 (21.7)	
	Unable to move	51 (6.9)	204,744 (2.3)
Lifestyle	Moderate to Vigorous Exercise	374 (50.3)	1,764,305 (20.7)
	Some physical activity	252 (33.9)	2,083,695 (24.5)
	No physical activity	117 (15.7)	4,652,856 (54.7)
	No alcohol use	385 (51.8)	2,226,486 (26.3)
	14 or less drinks per week	344 (46.3)	1,242,238 (14.6)
	>14 drinks per week	14 (1.9)	5,009,245 (59)
	Non-Smokers		7,422,711 (84.4)
	Never smoked	272 (36.6)	
	Quit	404 (54.4)	
	Smokers	67 (9.0)	1,372,885 (15.6)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

The purpose of this national cross-sectional survey was to describe aging with MS in Canada that would provide the data necessary to determine, in future analysis, the influence of personal, health and environmental factors on quality of life and disability among older people with MS. Incidence and prevalence estimates of MS in Canada are amongst the highest worldwide [34] and a shift in peak age prevalence of MS to older ages suggesting that persons with MS in Canada are living longer [2]. Interventions to promote successful aging with MS may take on more importance over the next decade since quality-of-life and productivity in advanced age is a desired outcome for most [6]. To our knowledge this is the most comprehensive study focusing on examination of health and lifestyle factors specifically among older persons with MS.

The prevalence of MS in Canada has been estimated to be 240 per 100,000 [34]. The CCHS estimates that there are 93,535 Canadians living with MS [31] and since Statistics Canada estimates that people over 55 make up about 27.7% of the population [35], there could be approximately 25,000 people with MS over 55 years of age. We attempted to contact as many older people with MS as possible and our sample (n=743) likely represents about 3% of this group, less than our initial 5% target. Data quality was very good with only 5.4% missing data and 709 respondents providing optional qualitative data. The high rates of response and completion and may be due to pre-screening of potential respondents at clinic sites. Only 7.5% of respondents required assistance by the research team but we were unable to discern survey completion by proxy.

With an average age about 65 years, our respondents are 10 to 25 years older than cohorts examined in previous studies of older people with MS [36-42]. Compared to these previous

1
2
3 studies, which reported the number of years since MS diagnosis ranging from 8-19 years [37 38],
4
5 our participants have been diagnosed with MS for 24.8 (± 10.0) years. Our respondents' level of
6
7 education aligns with most other studies of people with MS [37-39 41]. The group is fairly well-
8
9 educated which is not surprising considering characteristics of people who respond to surveys in
10
11 general [43]. When compared to the Canadian population [31], fewer of our respondents had
12
13 advanced education including post-secondary degrees and post-graduate degrees. Furthermore,
14
15 only about 10% of our sample was employed in any capacity which is substantially lower than
16
17 other studies of older people with MS [36 38 41]. Since maintaining employment is associated
18
19 with better quality-of-life, at least in younger people with MS [41], factors associated with
20
21 employment and retirement as one ages with MS will be an important area of future study.
22
23
24
25
26

27 More than a quarter of our cohort were severely disabled (28.6%); over ten times that of
28
29 aging counterparts in Canada (2.3%). They either were restricted to ambulation with a
30
31 wheelchair or walking with physical assistance indoors or were bedridden; an interesting finding
32
33 given that most respondents lived in private residences (96.2%) without home care, equipment or
34
35 home modifications. Since only about half of respondents had private health insurance, it is
36
37 possible that most people manage at home with few resources outside of the publicly-funded
38
39 health system and the care provided by spouses and children. Since the proportion of respondents
40
41 living in assisted-living and long-term care in this study was small, our sample may not be
42
43 representative of older people with MS living in these settings.
44
45
46
47

48 In terms of lifestyle behaviours, the participants in this study were less physically active
49
50 but less likely to drink alcohol or smoke compared to their Canadian counterparts [31]. In large
51
52 cohorts of people with MS in the United Kingdom (n=895) and Belgium (n=1372), healthy
53
54 lifestyle behaviours are associated with less severe and less rapid disability progression [44 45].
55
56
57
58
59
60

Poor lifestyle behaviors, specifically smoking, have been associated with increased risk of comorbid diseases [46]. Future studies will examine the role of health behaviors and disability in this group of older people with MS.

Although this unique cohort may provide new insights on aging with MS, there are some study limitations. The cross sectional design limits our ability to assess change and the effects of variables on individual outcomes over time. By nature of the volunteer survey design, our sample may be biased in that active participants were more likely to respond. Even though we provided assistance over the phone, those with significant disability who required assistance may have declined. The low sampling of persons in nursing homes and assisted living situations highlights the need for specialized recruitment strategies to study this sub-set of people living with MS.

Conclusion

Despite barriers to participation, we were able to recruit and gather detailed responses (with good data quality) from a large proportion of older Canadians with MS. Even with extensive pretesting, national collaboration, exhaustive sampling and high return rates, our recruitment (n=743) was less than our target (n=1250). However, we were able to recruit a large proportion of people who were severely disabled even though only 7.5% of respondents requested assistance. The data suggests that this sample of older people with MS are less likely to be employed, are less active and more disabled than other older Canadians.

Acknowledgements

This study was funded by a Canadian Institutes of Health Research (CIHR) Post-doctoral Fellowship for M.P., a Newfoundland and Labrador Centre for Applied Health Research,

Healthy Aging Research Program Project Grant, the Health Care Foundation (Eastern Health Authority) Research Project Grant, and the Physiotherapy Foundation of Canada Alberta Research Award. Support was also received from the MS Society of Canada and local chapters; provincial MS clinic neurologists, MS nurses, administrative and research staff; and the Rehabilitation and Continuing Care Program, Eastern Health Authority. We specifically acknowledge the work of site coordinators: Trudy Campbell, Heather Rosehart, Amy Cuthbertson, Darren Nickel, Pam Dumont, Madonna de Lemos, Deborah Goldberg, Irene Yee, Nicholas Hall, Maureen Perera, Sarah Williamson, Carolina Moriello and Vanessa Bouchard.

Contributorship

Concept development and funding applications- MP, JDF, MG
Ethics applications and participant recruitment-MP, CH, OJM, SH, JDF, ADS, PO, SAM, LMM, KK, PS, NM, RAM, MS
Data collection, management and analysis-MP, OJM, SB, CH, SH, PWA
Manuscript preparation and /or review-MP, OJM, CH, JDF, ADS, PO, SAM, LMM, PS, NM, RAM, MS, MG

Data sharing

Analysis on this study data is ongoing. Currently six additional unpublished manuscripts are in preparation.

Competing Interests

None

References

1. Sadovnick AD. Genetic background of multiple sclerosis. *Autoimmunity reviews* 2012;**11**(3):163-6.

2. Marrie RA, Yu N, Blanchard J, et al. The rising prevalence and changing age distribution of multiple sclerosis in Manitoba. *Neurology* 2010;**74**(6):465-71.

3. D'Hooghe M B, Haentjens P, Nagels G, et al. Alcohol, coffee, fish, smoking and disease progression in multiple sclerosis. *Eur J Neurol* 2012;**19**(4):616-24.

4. Marrie RA, Rudick R, Horwitz R, et al. Vascular comorbidity is associated with more rapid disability progression in multiple sclerosis. *Neurology* 2010;**74**(13):1041-7.

5. Ploughman M, Austin MW, Murdoch M, et al. Factors influencing healthy aging with multiple sclerosis: a qualitative study. *Disability and rehabilitation* 2012;**34**(1):26-33.

6. Ploughman M, Austin MW, Murdoch M, et al. The path to self-management: a qualitative study involving older people with multiple sclerosis. *Physiother Can* 2012;**64**(1):6-17.

7. He D, Guo R, Zhang F, et al. Rituximab for relapsing-remitting multiple sclerosis. *The Cochrane database of systematic reviews* 2013;**12**:CD009130.

8. Rietberg MB, Brooks D, Uitdehaag BM, et al. Exercise therapy for multiple sclerosis. *The Cochrane database of systematic reviews* 2005(1):CD003980.

9. Ploughman M, Austin M, Stefanelli M, et al. Applying cognitive debriefing to pre-test patient-reported outcomes in older people with multiple sclerosis. *Qual Life Res* 2010;**19**(4):483-7.

10. Canada MSo. MS Research Portal. Secondary MS Research Portal 2012. <http://msresearch.ca/> (accessed May 2014).

11. Hobart J, Lamping D, Fitzpatrick R, et al. The Multiple Sclerosis Impact Scale (MSIS-29): a new patient-based outcome measure. *Brain* 2001;**124**(Pt 5):962-73.

12. Mahoney FI, Barthel DW. Functional Evaluation: The Barthel Index. *Md State Med J* 1965;**14**:61-5.

13. Nicholl L, Hobart J, Dunwoody L, et al. Measuring disability in multiple sclerosis: is the Community Dependency Index an improvement on the Barthel Index? *Mult Scler* 2004;**10**(4):447-50.

14. Cano SJ, O'Connor RJ, Thompson AJ, et al. Exploring disability rating scale responsiveness II: do more response options help? *Neurology* 2006;**67**(11):2056-9.

15. van der Putten JJ, Hobart JC, Freeman JA, et al. Measuring change in disability after inpatient rehabilitation: comparison of the responsiveness of the Barthel index and the Functional Independence Measure. *J Neurol Neurosurg Psychiatry* 1999;**66**(4):480-4

16. Yeo D, Faleiro, R., Lincoln, N.B. Barthel. ADL Index: a comparison of administration methods. *Clin Rehabil* 1995;**9**(1):34-9.
17. Marrie RA, Horwitz RI. Emerging effects of comorbidities on multiple sclerosis. *Lancet Neurol* 2010;**9**(8):820-8.
18. Horton M, Rudick RA, Hara-Cleaver C, et al. Validation of a self-report comorbidity questionnaire for multiple sclerosis. *Neuroepidemiology* 2010;**35**(2):83-90.
19. Wade DT, Legh-Smith J, Langton Hewer R. Social activities after stroke: measurement and natural history using the Frenchay Activities Index. *Int Rehabil Med* 1985;**7**(4):176-81.
20. Turnbull JC, Kersten P, Habib M, et al. Validation of the Frenchay Activities Index in a general population aged 16 years and older. *Arch Phys Med Rehabil* 2000;**81**(8):1034-8.
21. Carter J, Mant F, Mant J, et al. Comparison of postal version of the Frenchay Activities Index with interviewer-administered version for use in people with stroke. *Clin Rehabil* 1997;**11**(2):131-8.
22. Einarsson U, Gottberg K, Fredrikson S, et al. Activities of daily living and social activities in people with multiple sclerosis in Stockholm County. *Clin Rehabil* 2006;**20**(6):543-51.
23. Ytterberg C, Johansson S, Gottberg K, et al. Perceived needs and satisfaction with care in people with multiple sclerosis: a two-year prospective study. *BMC Neurol* 2008;**8**:36.
24. Godwin M, Streight S, Dyachuk E, et al. Testing the Simple Lifestyle Indicator Questionnaire: Initial psychometric study. *Can Fam Physician* 2008;**54**(1):76-7.
25. Honarmand K, Feinstein A. Validation of the Hospital Anxiety and Depression Scale for use with multiple sclerosis patients. *Mult Scler* 2009;**15**(12):1518-24.
26. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;**67**(6):361-70.
27. Wagnild GM, Young HM. Development and psychometric evaluation of the Resilience Scale. *J Nurs Meas* 1993;**1**(2):165-78.
28. Weinert C. A social support measure: PRQ85. *Nurs Res* 1987;**36**(5):273-7.
29. Weinert C, Cudney S, Hill WG. Rural women, technology, and self-management of chronic illness. *Can J Nurs Res* 2008;**40**(3):114-34.
30. Long KA, Weinert C. Descriptions and perceptions of health among rural and urban adults with multiple sclerosis. *Res Nurs Health* 1992;**15**(5):335-42.
31. Canada S. Canadian Community Health Survey: Data Liberation Initiative 2010. Available from: equinox2.uwo.ca (accessed May 2014).
32. Gagne L, Tedds, L., Sullivan, J., Berniaz, K. An introduction to the UT/DLS: Microdata analysis subsetting. Secondary An introduction to the UT/DLS: Microdata analysis

subsetting n.d. <http://sda.chass.utoronto.ca/sdaweb/doc/SDAtutorial2008.pdf> (accessed May 2014).

33. Hawthorne G, Elliott P. Imputing cross-sectional missing data: comparison of common techniques. *Aust N Z J Psychiatry* 2005;**39**(7):583-90.

34. Evans C, Beland SG, Kulaga S, et al. Incidence and prevalence of multiple sclerosis in the Americas: a systematic review. *Neuroepidemiology* 2013;**40**(3):195-210.

35. Canada S. Population by sex and age group Secondary Population by sex and age group 2012. <http://www.statcan.gc.ca/tables-tableaux/sum-som/l01/cst01/demo10a-eng.htm> (accessed May 2014).

36. Buchanan RJ, Chakravorty BJ, Tyry T, et al. Age-related comparisons of people with multiple sclerosis: demographic, disease, and treatment characteristics. *NeuroRehabilitation* 2009;**25**(4):271-8.

37. Giordano A, Ferrari G, Radice D, et al. Self-assessed health status changes in a community cohort of people with multiple sclerosis: 11 years of follow-up. *Eur J Neurol* 2013;**20**(4):681-8.

38. Asano M, Duquette P, Andersen R, et al. Exercise barriers and preferences among women and men with multiple sclerosis. *Disability and rehabilitation* 2013;**35**(5):353-61.

39. Kasser SL, Kosma M. Health beliefs and physical activity behavior in adults with multiple sclerosis. *Disabil Health J* 2012;**5**(4):261-8.

40. Jensen MP, Molton IR, Gertz KJ, et al. Physical activity and depression in middle and older-aged adults with multiple sclerosis. *Disabil Health J* 2012;**5**(4):269-76.

41. Krokavcova M, Nagyova I, van Dijk JP, et al. Mastery, functional disability and perceived health status in patients with multiple sclerosis. *Eur J Neurol* 2008;**15**(11):1237-44.

42. Kayes NM, McPherson KM, Schluter P, et al. Exploring the facilitators and barriers to engagement in physical activity for people with multiple sclerosis. *Disability and rehabilitation* 2011;**33**(12):1043-53.

43. Chretien JP, Chu LK, Smith TC, et al. Demographic and occupational predictors of early response to a mailed invitation to enroll in a longitudinal health study. *BMC Med Res Methodol* 2007;**7**:6.

44. Manouchehrinia A, Tench CR, Maxted J, et al. Tobacco smoking and disability progression in multiple sclerosis: United Kingdom cohort study. *Brain* 2013;**136**(Pt 7):2298-304.

45. D'Hooghe M B, Nagels G, De Keyser J, et al. Self-reported health promotion and disability progression in multiple sclerosis. *J Neurol Sci* 2013;**325**(1-2):120-6.

46. Marrie RA, Horwitz RI, Cutter G, et al. Smokers with multiple sclerosis are more likely to report comorbid autoimmune diseases. *Neuroepidemiology* 2011;**36**(2):85-90.




Figure 1: Survey Respondents
121x101mm (150 x 150 DPI)




Figure 2: Number of respondents mapped by postal code
99x58mm (300 x 300 DPI)

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Complete
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	✓
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	✓
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	✓
Objectives	3	State specific objectives, including any prespecified hypotheses	✓
Methods			
Study design	4	Present key elements of study design early in the paper	✓
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	✓
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up <i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	✓
		(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed <i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	✓
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	✓
Bias	9	Describe any efforts to address potential sources of bias	✓
Study size	10	Explain how the study size was arrived at	✓
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	✓
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	✓
		(b) Describe any methods used to examine subgroups and interactions	✓
		(c) Explain how missing data were addressed	✓
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	✓
		(e) Describe any sensitivity analyses	✓

Continued on next page

Results

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	✓
		(b) Give reasons for non-participation at each stage	✓
		(c) Consider use of a flow diagram	
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	✓
		(b) Indicate number of participants with missing data for each variable of interest	✓
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	✓
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	✓
		(b) Report category boundaries when continuous variables were categorized	✓
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	✓
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	✓

Discussion

Key results	18	Summarise key results with reference to study objectives	✓
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	✓
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	✓
Generalisability	21	Discuss the generalisability (external validity) of the study results	✓

Other information

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	✓
---------	----	---	---

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

The Canadian Survey of Health, Lifestyle and Aging with Multiple Sclerosis; Methodology and Initial Results

Journal:	<i>BMJ Open</i>
Manuscript ID:	bmjopen-2014-005718.R1
Article Type:	Research
Date Submitted by the Author:	12-Jun-2014
Complete List of Authors:	Ploughman, Michelle; Memorial University of Newfoundland, Faculty of Medicine Beaulieu, Serge; Eastern Health Authority, Harris, Chelsea; Memorial University, Faculty of Medicine Hogan, Stephen; Memorial University, Faculty of Medicine Manning, Olivia; Memorial University, Faculty of Medicine Alderdice, Penelope; Memorial University, Faculty of Medicine Fisk, John; Dalhousie University, Sadovnick, A; University of British Columbia, Department of Medical Genetics and Division of Neurology O'Connor, Paul; St. Michaels Regional Hospital, Department of Neurology Morrow, Sarah; London Health Science Centre, Metz, Luanne; University of Alberta, Department of Neurology Smyth, Penelope; University of Calgary, Clinical Neurosciences Mayo, Nancy; McGill University, Medicine Marrie, Ruth; University of Manitoba, Departments of Internal Medicine and Community Health Sciences Knox, Katherine; University of Saskatchewan, Department of Physical Medicine and Rehabilitation Stefanelli, Mark; Memorial University, Faculty of Medicine Godwin, Marshall; Memorial University, Primary Health Care Research Unit
Primary Subject Heading:	Neurology
Secondary Subject Heading:	Research methods
Keywords:	Multiple sclerosis < NEUROLOGY, REHABILITATION MEDICINE, NEUROLOGY

SCHOLARONE™
Manuscripts

The Canadian Survey of Health, Lifestyle and Aging with Multiple Sclerosis; Methodology and Initial Results

Michelle Ploughman PT, PhD*, Serge Beaulieu**, Chelsea Harris BKin MPH*, Stephen Hogan BKin*, Olivia J. Manning*, Penelope W. Alderdice PhD*, John D. Fisk PhD[‡], A. Dossa Sadovnick PhD[‡], Paul O'Connor, MD, FRCPC[‡], Sarah A. Morrow, MD[‡], FRCPC, Luanne M. Metz MD, FRCPC[#], Penelope Smyth MD, FRCPC[‡], Nancy Mayo, PhD^{§§}, Ruth Ann Marrie, MD, PhD^{††}, Katherine B. Knox, MD, FRCPC^{‡‡}, Mark Stefanelli MD FRCPC^{¶¶}, Dr. Marshall Godwin, MD^{##}

*Recovery and Performance Laboratory, Rehabilitation Research Unit, Faculty of Medicine, Memorial University, St. John's NL Canada

**Eastern Health Authority, St. John's NL Canada

‡Dalhousie University, Halifax NS, Canada

†Department of Medical Genetics and Division of Neurology, Faculty of Medicine University of British Columbia, Vancouver BC, Canada

‡Department of Neurology, St. Michaels Regional Hospital, Toronto ON, Canada

¶London Health Science Centre, London ON, Canada

‡Department of Neurology, University of Alberta, Edmonton AB, Canada

‡Clinical Neurosciences, University of Calgary, Calgary AB, Canada

§§Clinical Epidemiology, McGill University, Montreal QC, Canada

††Departments of Internal Medicine and Community Health Sciences, University of Manitoba, Winnipeg MB, Canada

‡‡Department of Physical Medicine and Rehabilitation, College of Medicine, University of Saskatchewan, Saskatoon SK, Canada

¶¶Neurology, Faculty of Medicine, Memorial University, St. John's NL Canada

##Primary Health Care Research Unit, Faculty of Medicine, Memorial University, St. John's NL, Canada

Corresponding Author/Research Site

Dr. Michelle Ploughman, Recovery and Performance Laboratory, Rehabilitation Research Unit, Faculty of Medicine, Memorial University, Rm 400, L.A. Miller Centre, 100 Forest Rd., St. John's NL Canada. Ph 709 777 2099; fax 709 777 8702; email mploughm@mun.ca

Key Words: aging; diet; exercise; health behaviours; multiple sclerosis; neurodegenerative disease; older persons; quality of life

Short Title: Health Lifestyle and Aging with Multiple Sclerosis

Tables: 3

Figures: 4

For peer review only

ABSTRACT

Objective

People with multiple sclerosis (MS) are living longer so strategies to enhance long term health and quality of life are garnering more interest. We aimed to create a profile of aging with MS in Canada by recruiting 1250 (5% of the Canadian population over 55 with MS) participants and focusing data collection on health and lifestyle factors, disability, participation and quality of life in order to determine factors associated with healthy aging.

Design

National multi-centre postal survey

Setting

Recruitment from Canadian MS clinics, MS Society of Canada chapters, and newspaper advertisements

Participants

People aged 55 years or older with MS symptoms more than 20 years

Outcome Measures

Validated outcome measures and custom-designed questions examining MS disease characteristics, living situation, disability, co-morbid conditions, fatigue, health behaviours, mental health, social support, impact of MS, and others.

Results

Of the 921 surveys, 743 were returned (80.7% response rate). Participants (mean age 64.6 ±6.2 years) reported living with MS symptoms for an average of 32.9±9.5 years and 28.6% were either wheelchair users or bedridden. There was only 5.4% missing data and 709 respondents provided optional qualitative information. According to data derived from the 2012 Canadian Community Health Survey of Canadians over 55 years of age, older people with MS from this survey sample are about 8 times less likely to be employed full-time. Older people with MS were less likely to engage in regular physical activity (26.7%) compared to typical older Canadians (45.2%). However, they were more likely to abstain from alcohol and smoking.

Conclusion

Despite barriers to participation, we were able to recruit and gather detailed responses (with good data quality) from a large proportion of older Canadians with MS. The data suggests that this sample of older people with MS are less likely to be employed, are less active and more disabled than other older Canadians.

Strengths and limitations of this study

- This unique cohort provides new insights on aging with MS. With an average age about 65 years, our respondents are 10 to 25 years older than cohorts examined in previous studies of older people with MS.
- Our recruitment methods and a very high survey return rate allowed us to obtain data from about 3% of our target population.
- With only about 10% of our sample being employed in any capacity, our results point to a need to understand retirement and employment in aging with MS.
- The cross sectional design limits our ability to assess change and the effects of variables on individual outcomes over time.
- The low sampling of persons in nursing homes and assisted living situations highlights the need for specialized recruitment strategies to study this sub-set of people living with MS.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Multiple Sclerosis (MS) is a highly unpredictable and heterogeneous autoimmune disease diagnosed between the ages of 18-40 years. Affecting women more than men, the cause of MS is unknown, however research suggests that a combination of environmental factors and genetic predisposition may be the trigger for disease onset [1]. The number of people with MS over the age of 55 is increasing, possibly due to improved longevity and more effective treatments [2]. On the other hand, several authors have shown that smoking [3] and cardiovascular disease [4] accelerate disease progression. People with MS are interested in strategies that minimize disability and enhance role participation and quality of life [5]. The uncertainty surrounding MS progression along with aging MS patients warrants the need for investigation into potential disease modifying health behaviours. Older individuals who have lived with MS for many years may provide insight into factors that contribute to living a long and healthy life with MS [6].

Older people with MS are often excluded from MS research. Studies involving clinical drug trials [7] and exercise interventions [8], in particular, target patients with mild relapsing-remitting disease. Furthermore, because they may not require intensive medical oversight associated with taking disease-modifying drugs, older patients may not attend MS clinics as often as those recently diagnosed. The older MS cohort are also more likely to be disabled and living in an institutional setting, making recruitment challenging. In addition, disability-related impairments such as those affecting vision, cognition and coordination complicate survey and patient-reported outcome administration [9].

1
2
3 In order to create a profile of aging with MS in Canada, we developed a national cross-
4 sectional survey, based on previous qualitative research [5 6], using validated patient-reported
5 outcome measures [9]. We focused on health and lifestyle factors, disability, participation and
6 quality of life, targeting people 55 years of age and older with MS symptoms for 20 or more
7 years. By partnering with major MS clinic registries and MS society chapters, we hypothesized
8 that we would be able to reach a representative sample of about 5% of the 'aging with MS in
9 Canada' cohort.
10
11
12
13
14
15
16
17
18
19

20 21 22 METHODS

23
24 **Participants and Sampling:** We targeted Canadians, aged 55 years or older, who had had self-
25 reported MS symptoms for 20 or more years. Prior to recruitment, the study was approved by 11
26 health research ethics boards in eight Canadian provinces. Between May 2011 and December
27 2012, participants were recruited by telephone from lists of eligible MS clinic patients and by
28 public advertising through the MS Society of Canada Research Internet Portal [10], local MS
29 Society chapters and their newsletters, and newspaper advertisements. A confirmed diagnosis
30 from a neurologist was available for participants recruited through the MS clinic databases; self-
31 identified participants provided a diagnosis date and neurologist name to confirm their diagnosis.
32 Rather than calculating a sample size, we used an exhaustive sampling technique to obtain a
33 large and representative sample; for example repeatedly targeting regional newspapers,
34 following-up at study sites, making formal and informal contacts with MS Society Chapter
35 support groups.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

52
53 After obtaining consent using a standard script, we mailed participants the questionnaire,
54 with an information sheet, a toll-free telephone helpline number and email address and a prepaid
55
56
57
58
59
60

reply envelope. We requested that participants return the completed survey in 2 weeks; non-responders were telephoned for follow-up at 4 weeks. Participants who wished to complete the survey by telephone were encouraged to do so. Each questionnaire was coded to maintain anonymity.

Survey Design: The questionnaire consisted of validated patient-reported outcome tools and additional custom-designed questions chosen based on pilot research in a small sample of older people with MS [9]. We determined during this step that participants preferred a paper rather than online version of the survey as well as larger 14-point font, divided into sections that grouped similar questions together. Respondents were instructed to complete the survey in these sections (chapters) taking a break in between. Since we previously observed confusion with questions involving temporal parameters, questions regarding activity involvement that followed the same timeline (e.g. weekly, monthly, or yearly) were placed in sequence. To further examine the acceptability of the survey and to minimize skipped or misunderstood items, we asked the first 14 participants (11 returns) to provide feedback on the content, format and response burden. This feedback was integrated into the final version of the questionnaire but the responses were not included in the final data.

In order to include people with MS in the province of Quebec, Canada, the entire survey including the cover letter was translated into french by one independent professional translator. We also obtained french versions of previously validated tools. Four native french speakers involved in MS research made corrections to the initial translation in sequence. The corrections were then consolidated by a fifth french speaking MS researcher. The advertisement and

consent scripts were translated by a native french speaking MS researcher and reviewed and corrected by a second.

Survey Content: The final survey consisted of five chapters (**Table 1**) ending with open response questions for participants to provide additional comments. Where appropriate, we obtained permission from the developers to use previously validated tools.

We used the Multiple Sclerosis Impact Scale (MSIS-29), the Barthel Index and the Co-morbidity Questionnaire to assess health. The MSIS-29 consists of 29 questions: 20 address the physical impact and nine address the psychological impact of MS generating separate or combined scores as a percentage of 100. Higher scores indicate greater impact of MS [11]. The MSIS-29 physical component correlates with the Barthel Index (0.71) and the psychological component with EQ-5D (0.68). It is reported to have good responsiveness with large to moderate effect sizes [12]. The Barthel Index measures function and mobility in activities of daily living (ADL) in ten areas: feeding, bathing, grooming, dressing, bowel control, bladder control, toileting, transfers, ambulation and stair climbing [13]. Items are rated in terms of whether individuals can perform activities independently, with some assistance, or are completely dependent, with a total score ranging from 0 (complete assistance required) to 100 (independent). The Barthel Index has been tested in people with MS [14-16] and is reliable whether administered by telephone, in-person or by mail [17] with effect sizes comparable to the Functional Independence Measure (0.34) [18]. The list of co-morbidities used in this survey was adapted from Marrie and Horwitz [19 20] in consultation with Dr. Marrie (personal communication).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

We determined the degree of participation in instrumental activities of daily living (iADL) using the Frenchay Activities Index (FAI) and individual components of lifestyle (exercise, diet, smoking, alcohol, and stress) using the Simple Lifestyle Indicator Questionnaire (SLIQ). The FAI assesses the frequency of participation in household chores, social activities, work and leisure for a summed raw score of 15 to 60; higher score indicating more frequent participation [21]. The FAI has excellent test-retest reliability as a mailed questionnaire (0.96) [22] and correlates well with face-to-face interview scores [23]. The FAI has been mainly used in stroke but has also been used in large MS population studies [24 25]. The SLIQ was developed to measure the individual components of lifestyle that affect cardiovascular disease [26]. Initial content and face validity were assessed by the developers among 136 people attending a family practice clinic[26] and SLIQ scores were highly correlated with those of blinded reviewers (0.77). We were also interested in mental health, coping and social support in aging with MS. The Hospital Anxiety and Depression Scale (HADS) is a self-assessment scale found to reliably screen for depression and anxiety in multiple sclerosis patients [27]. It is divided into an Anxiety subscale (HADS-A) and a Depression subscale (HADS-D) each containing seven intermingled items scored out of 21; higher score indicating greater symptoms [28]. A threshold score of 8 or higher on HADS subscales had high sensitivity (Depression 90%, Anxiety 88.5%) and specificity (Depression 87.3%, Anxiety 80.7%) compared to structured interview using DSM-1 criteria among people with MS [27]. Resilience, the ability to overcome adversity, was measured using the 25 item Resilience Scale. Responses to the test questions range from 1 (disagree) to 7 (agree) and the summed score can be further interpreted as, low resilience (<120), moderate resilience (125-145), and high resilience (>145) [29] and has been validated in older community-

living individuals with high internal consistency reliability (0.91)[29] . The Personal Resource Questionnaire (PRQ-2000) measures social support and consists of 15 statements with which the subject agrees or disagrees based on a seven-point Likert scale; higher scores indicating higher support (e.g. I belong to a group in which I feel important 1-Strongly Disagree to 7-Strongly Agree). PRQ-2000 demonstrates good internal consistency (0.87 to 0.93), adequate construct validity (moderate correlations with anxiety -0.37, neuroticism -0.28 and extroversion 0.32) and good predictive and construct validity in multiple chronic disease data sets [30]. Mailed versions have been used in community-living women with chronic health conditions [31] and adults with MS [32].

Table 1: Survey Content

CHAPTER	DOMAINS	MEASUREMENT TOOLS
You and Your MS (5 pages)	Demographics	Postal code, gender, height, weight, age, finances and education
	MS Disease Characteristics	Symptom onset, time since diagnosis, and type of MS
	Environment	Living situation, accessibility and home modifications
Your Health (6 pages)	Health-Related Quality of Life	Visual analogue scale of perceived health status [9] Multiple Sclerosis Impact Scale-29 [11]
	Physical Health and Disability	Barthel Index [12] Co-Morbidity Questionnaire [17]
	Health Care Services	Use and satisfaction with health care services, medications and complementary/alternative health.
Your Activities (7 pages)	Participation and instrumental Activities of Daily Living	Frenchay Activities index [19]; Employment and impact of fatigue
	Lifestyle	Simple Lifestyle Indicator Questionnaire (diet, exercise, stress, smoking, alcohol) [24]
Your Thoughts and Feelings (8 pages)	Mental/ Cognitive Health	Hospital Anxiety and Depression Scale (HADS) Resilience Scale [25] Problems with concentration and memory
Your Social Life (4 pages)	Social Support	Personal Resource Questionnaire-2000 [30]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Canadian comparison data: Comparative data was collected from the Canadian Community Health Survey (CCHS) 2012 version; a national cross-sectional survey that gathers information about health, social and economic status and lifestyle behaviours of the population. Data was extracted from the CCHS database using the University of Toronto Data Library Service (DLS) [33]. A CCHS sample was selected based on age (>55 years). Data was collected from CCHS questions that matched question from our survey (e.g. mobility from the Barthel Index and lifestyle from the SLIQ). Outputs (ie. highest level of education, employment status) were analyzed using Survey Documentation & Analysis (SDA) software [34]. Frequency tables were created for variables of interest.

Data Cleaning and Imputation: We custom designed the database application to house project data and minimize error during data imputation. Missing data was imputed using three techniques: (1) deductive imputation, (2) person mean substitution, and (3) ‘hot deck’ imputation [35]. The deductive technique was used to impute missing data when there was sufficient comments and answers to related questions. If less than half of the data was missing on a section, scale, or subscale, person mean substitution (PMS) was used. PMS takes the mean score of the particular participant on that section, scale, or subscale and uses the mean as the value for the missing data. If greater than half of the data was missing on a section, scale, or subscale, ‘hot deck’ imputation was used [35]. This involved taking a value from the "hot deck" or in other terms, finding another participant of the same gender, age (± 5 years), education level, and living in the same province. Hot deck imputation was selected for this study because it is proven to be the most efficient method of data imputation as it is more robust in large samples when

1
2
3 compared to other imputation methods [35]. The exact scores for each question from the
4
5 substitute participant were imputed for the missing values for the participant of interest.
6
7
8 Variables that could not be imputed and remained empty included, from **Table 1**, Demographics
9
10 (e.g. weight, height, finances), MS Disease Characteristics, Environment, and Health Care
11
12 Services.
13

14
15
16
17 **Data Analysis:** Data was analyzed using IBM SPSS software v20. Frequencies and descriptive
18
19 statistics (mean, SD and data range) were used to determine subject characteristics.
20
21
22
23
24
25
26

27 RESULTS

28 *Survey Response*

29
30
31 We aimed to recruit 1250 participants for the study and through exhaustive sampling we
32
33 contacted 921 (74% of target) people with MS who met inclusion criteria. Seven hundred and
34
35 forty three participants returned the survey (**Figure 1**; 80.7% response rate). Survey response by
36
37 province is illustrated in **Figure 2**. Of the respondents, 90.4% (n=672) were identified through
38
39 provincial MS clinics while the remaining 71 self-identified by responding to newspaper ads and
40
41 word of mouth. Forty-five people requested help to complete the survey, and 11 participants
42
43 asked that the entire survey be completed by phone due to writing or vision difficulties or both.
44
45
46 Additional content was provided by 709 participants in the open-ended questions at the end of
47
48 the survey.
49
50
51

52
53 Only 563 of 104,763 cells contained missing data (5.37%). Almost every survey
54
55 contained one or two random missing items; in some cases answers to these could be inferred
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

using deductive reasoning (67/104,763=0.06%). We observed no pattern of missing data other than body weight (which we did not impute). The PMS technique was required for 170 cells (170/94,763=0.16%) and the ‘hot deck’ technique was required in 14 surveys (324 cells/104,763=0.31%). Cases of consecutive missing data seemed to be mainly due to pages adhering together resulting in questions being overlooked.

Insert Figure 1 about here

Insert Figure 2 about here

Sample Characteristics

Participants ranged in age from 55 to 88 years with a mean age of 64.6 years (±6.2). They described experiencing MS symptoms for 20 to 73 years with a mean duration of 32.9 years (±9.5). The respondents reported that the average time from onset of symptoms to diagnosis by a neurologist was 8.1 years (± 9.4) with a range from 0 to 47 years. The number of women outnumbered men 3.5:1 (**Table 2**).

Sixty percent reported at least one year of post-secondary education, and 90% were either retired or unemployed. About twenty-eight percent of the sample (n=212) reported that they were no longer able to walk and either used a wheelchair or were bedridden. The characteristics of the sample are fully described in **Table 2**.

Respondents were categorized by level of disability based on results of the Barthel Index (**Figure 3**) [18]. We measured activity and participation using the FAI which is divided into household (domestic) activity, leisure activity and outdoor activity (**Figure 4**). Nearly half

(50.6%) of the sample frequently participated in household activities but only 13.5% and 27.9% reported that they frequently engaged in leisure activities and outdoor activities respectively.

Table 2: Sample Characteristics

Characteristic		N (%)
Years with MS Symptoms	20-29	324 (43.6)
	30-39	245 (33.0)
	40-49	135 (18.2)
	50-59	31 (4.2)
	60 or more	7 (0.9)
Initial MS Diagnosis	Relapsing-Remitting	386 (52.0)
	Primary Progressive	99 (13.3)
	Secondary Progressive	65 (8.7)
	Progressive-Relapsing	16 (2.2)
	Benign	43 (5.8)
	Unknown	126 (17.0)
Living situation	Living in a private home	715 (96.2)
	Living in an assisted living complex	16 (2.2)
	Living in long term care	11 (1.5)
	Home Care	146 (19.7)
	Equipment	200 (26.9)
	Home Modifications	73 (9.8)
Private Insurance (Health/Disability)	Yes	393 (52.9)
	No	350 (47.1)

Insert Figure 3 about here

Insert Figure 4 about here

Older people with MS compared to typical older Canadians

A comparison sample of older people (>55) in Canada was derived from the 2012 Canadian Community Health Survey (CCHS; **Table 3**). CCHS respondents over the age of 55 years primarily lived with their spouse or partner (52.5%), with their spouse or partner and children (11.3%), or alone (22.7%). Only approximately one percent of this sample were living

with their parents. According to data derived from the CCHS, older people with MS from our survey sample have less post-secondary education and are about 8 times less likely to be employed full-time. About half of the MS survey sample have major mobility challenges (unable to walk outdoors, use a wheelchair or are bedridden) compared to only 9% in the CCHS sample. Older people with MS were more likely to engage in physical activity (69.4%) compared to typical older Canadians (45.3%). They were also more likely to abstain from alcohol and smoking.

Table 3: Comparison of Sample Characteristics to Others Aging in Canada

Characteristic		Study sample (%)	CCHS sample (%)
Age	55-64	414 (55.7)	4,671,364.6 (48.9)
	65-74	274 (36.9)	2,891,671.1 (30.3)
	75+	55 (7.4)	1,989,358.6 (20.8)
Gender	Males	166 (22.3)	4,568,522.0 (47.8)
	Females	577 (77.7)	4,983,872.3 (52.2)
Education	Less than high school	74 (10.0)	1,241,946 (15.2)
	High School	222 (29.9)	1,083,618 (13.3)
	1-2 year post-secondary	189 (25.4)	4,088,252 (4.9)
	Complete Post-graduate	258 (34.7)	5,442,586 (66.6)
Employment Status	Retired	504 (67.8)	2,878,200 (46.3)
	Unemployed	166 (22.3)	302,942 (4.8)
	Employed Full-time	30 (4.0)	2,068,571 (33.3)
	Employed Part-time	19 (2.6)	698,125 (11.2)
	Employed Casual	24 (3.2)	266,419 (4.3)
Mobility	Walk independently with or without cane outdoors	351 (47.2)	8,004,425 (90.9)
	Walk with cane or help indoors	180 (24.2)	593,584 (6.7)
	Wheelchair	161 (21.7)	
	Unable to move	51 (6.9)	204,744 (2.3)
Lifestyle	Moderate to Vigorous Exercise	193 (25.9)	1,764,305 (20.7)
	Some physical activity	316 (42.5)	2,083,695 (24.5)
	No physical activity	234 (31.5)	4,652,856 (54.7)
	No alcohol use	385 (51.8)	2,226,486 (26.3)
	14 or less drinks per week	344 (46.3)	1,242,238 (14.6)
	>14 drinks per week	14 (1.9)	5,009,245 (59)
	Non-Smokers	676 (91.0)	7,422,711 (84.4)
	Never smoked	272 (36.6)	
	Quit	404 (54.4)	
	Smokers	67 (9.0)	1,372,885 (15.6)

DISCUSSION

The purpose of this national cross-sectional survey was to describe aging with MS in Canada that would provide the data necessary to determine, in future analysis, the influence of personal, health and environmental factors on quality of life and disability among older people with MS. Incidence and prevalence estimates of MS in Canada are amongst the highest worldwide [36] and a shift in peak age prevalence of MS to older ages suggesting that persons with MS in Canada are living longer [2]. Interventions to promote successful aging with MS may take on more importance over the next decade since quality-of-life and productivity in advanced age is a desired outcome for most [6]. To our knowledge this is the most comprehensive study focusing on examination of health and lifestyle factors specifically among older persons with MS.

The prevalence of MS in Canada has been estimated to be 240 per 100,000 [36]. The CCHS estimates that there are 93,535 Canadians living with MS [33] and since Statistics Canada estimates that people over 55 make up about 27.7% of the population [37], there could be approximately 25,000 people with MS over 55 years of age. We attempted to contact as many older people with MS as possible and our sample (n=743) likely represents about 3% of this group, less than our initial 5% target. Data quality was very good with only 5.4% missing data and 709 respondents providing optional qualitative data. The high rates of response and completion and may be due to pre-screening of potential respondents at clinic sites. Only 7.5% of respondents required assistance by the research team but we were unable to discern survey completion by proxy.

With an average age about 65 years, our respondents are 10 to 25 years older than cohorts examined in previous studies of older people with MS [38-44]. Compared to these previous

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

studies, which reported the number of years since MS diagnosis ranging from 8-19 years [39 40], our participants have been diagnosed with MS for 24.8 (\pm 10.0) years. Our respondents' level of education aligns with most other studies of people with MS [39-41 43] .The group is fairly well-educated which is not surprising considering characteristics of people who respond to surveys in general [45]. When compared to the Canadian population [33], fewer of our respondents had advanced education including post-secondary degrees and post-graduate degrees. Furthermore, only about 10% of our sample was employed in any capacity which is substantially lower than other studies of older people with MS [38 40 43]. Since maintaining employment is associated with better quality-of-life, at least in younger people with MS [43], factors associated with employment and retirement as one ages with MS will be an important area of future study.

More than a quarter of our cohort were severely disabled (28.6%); over ten times that of aging counterparts in Canada (2.3%). They either were restricted to ambulation with a wheelchair or walking with physical assistance indoors or were bedridden; an interesting finding given that most respondents lived in private residences (96.2%) without home care, equipment or home modifications. Since only about half of respondents had private health insurance, it is possible that most people manage at home with few resources outside of the publicly-funded health system and the care provided by spouses and children. It was interesting that only a small percentage of survey respondents engaged in leisure or outdoor activity (Figure 4). Since the proportion of respondents living in assisted-living and long-term care in this study was small, our sample may not be representative of older people with MS living in these settings.

In terms of lifestyle behaviours, the participants in this study were more physically active and less likely to drink alcohol or smoke compared to their Canadian counterparts [33]. In large cohorts of people with MS in the United Kingdom (n=895) and Belgium (n=1372), healthy

lifestyle behaviours are associated with less severe and less rapid disability progression [46 47]. Poor lifestyle behaviors, specifically smoking, have been associated with increased risk of comorbid diseases [48] and significantly lower HRQOL when compared to people with MS who have never smoked [49]. Future studies will examine the role of health behaviors and disability in this group of older people with MS.

Although this unique cohort may provide new insights on aging with MS, there are some study limitations. The cross sectional design limits our ability to assess change and the effects of variables on individual outcomes over time. By nature of the volunteer survey design, our sample may be biased in that active participants were more likely to respond. Even though we provided assistance over the phone, those with significant disability who required assistance may have declined. The low sampling of persons in nursing homes and assisted living situations and those older than 75 years highlights the need for specialized recruitment strategies to study this sub-set of people living with MS. Although preliminary testing indicated that participants preferred paper, a web-based version may have improved our recruitment. We also have compared our sample to the Canadian Census CCHS sample. The survey questions do not perfectly align so the values are estimates only. A final limitation of the survey was that we selected the Simple Lifestyle Indicator Questionnaire to assess health behaviours because of its short, simple design, so the data collected lacks detail on diet behaviours which will limit future analysis.

Conclusion

Despite barriers to participation, we were able to recruit and gather detailed responses (with good data quality) from a large proportion of older Canadians with MS. Even with extensive pretesting, national collaboration, exhaustive sampling and high return rates, our

recruitment (n=743) was less than our target (n=1250). However, we were able to recruit a large proportion of people who were severely disabled even though only 7.5% of respondents requested assistance. The data suggests that this sample of older people with MS are less likely to be employed, are more disabled but have more positive health behaviours than other older Canadians.

Acknowledgements

Support for this project was received from the MS Society of Canada and local chapters; provincial MS clinic neurologists, MS nurses, administrative and research staff; and the Rehabilitation and Continuing Care Program, Eastern Health Authority. We specifically acknowledge the work of site coordinators: Trudy Campbell, Heather Rosehart, Amy Cuthbertson, Darren Nickel, Pam Dumont, Madonna de Lemos, Deborah Goldberg, Irene Yee, Nicholas Hall, Maureen Perera, Sarah Williamson, Carolina Moriello and Vanessa Bouchard.

Contributorship Statement

Concept development and funding applications- MP, JDF, MG
Ethics applications and participant recruitment-MP, OJM, CH, SH, JDF, ADS, PO, SAM, LMM, PS, NM, RAM, MS
Data collection, management and analysis-MP, OJM, SB, CH, SH, PWA
Manuscript preparation and /or review-MP, OJM, CH, JDF, ADS, PO, SAM, LMM, PS, NM, RAM, MS, MG

Competing Interests

Dr. Michelle Ploughman was supported by a Canadian Institutes for Health Research (Institute of Aging) Postdoctoral Fellowship. Her research was supported by the Newfoundland and Labrador Centre for Applied Health Research, Health Care Foundation, and Physiotherapy Foundation of Canada.

Serge Beaulieu, Stephen Hogan, Dr. Penelope Alderdice, Dr. A. Dessa Sadovnick, Dr. Luanne M. Metz, and Dr. Nancy Mayo have no disclosures or conflicts of interest to report.

Chelsea Harris and Olivia Manning were supported by endMS Research and Training Network summer studentships.

Dr. John Fisk is the Director of the endMS Atlantic Regional Research and Training Centre, which is funded by the Multiple Sclerosis Society of Canada. He receives research funding from the Canadian Institutes of Health Research (CIHR) and in the past has received grants, honoraria and consultation fees from Astra- Zeneca, Bayer, Biogen-Idec Canada, Heron Evidence Development Limited, Hoffmann-La Roche, MAPI Research Trust, Novartis, Sanofi-Aventis, Serono Canada, and QualityMetric Inc.

Dr. Paul O'Connor has received consulting fees and/or research support for MS trials from Actelion, Bayer, Biogen Idec, BioMS, Cognosci, Daiichi Sankyo, EMD Serono, Genentech, Genmab, Novartis, Roche, Sanofi Aventis, Teva and Warburg Pincus.

Dr. Sarah Morrow has received research support from Biogen Idec and Novartis Canada, and received support for speaking engagement and advisory panels for Bayer, Biogen Idec, EMDSerono and Teva Neurosciences and Novartis Canada.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Dr. Penelope Smyth has served on scientific advisory boards for Novartis and EMD Serono. She has received funding for travel to pharmaceutically sponsored events by EMD Serono. She has received honoraria for speaking, facilitating and chairing from Novartis and EMD Serono as well as from the MS Society of Canada. She has received academic research support for medical education research from the Arnold P. Gold Foundation, and the Teaching and Learning Enhancement Fund from the University of Alberta.

Dr. Ruth Ann Marrie receives research funding from: Canadian Institutes of Health Research, Multiple Sclerosis Society of Canada, Multiple Sclerosis Scientific Research Foundation, Rx & D Health Research Foundation, and has conducted clinical trials funded by Sanofi-Aventis.

Dr. Katherine Knox received grant funding from Saskatoon City Hospital Foundation and MS Society of Canada, educational CME event grant funding from Bayer HealthCare Pharmaceuticals, honoraria /travel grant from Biogen and Teva. She was a member of the CIHR and MS Society of Canada Scientific Expert Working Group on CCSVI and MS.

Dr. Mark Stefanelli has served on advisory boards for Biogen Idec and Novartis and has been provided speaker/travel support by Teva Neuroscience. His research was supported by Genzyme and Biogen.

Dr. Marshall Godwin's research is funded by the Canadian Institutes of Health Research, Public Health Agency of Canada, Health Canada, and Newfoundland and Labrador Centre for Applied Health Research. He reports no conflicts of interest.

Funding

This research was supported by a Canadian Institutes of Health Research (CIHR) Post-doctoral Fellowship, a Newfoundland and Labrador Centre for Applied Health Research, Healthy Aging Research Program Project Grant, the Health Care Foundation (Eastern Health Authority) Research Project Grant, and the Physiotherapy Foundation of Canada Alberta Research Award.

Data Sharing

Additional data can be obtained by contacting Dr. Michelle Ploughman at
Michelle.Ploughman@med.mun.ca.

References

1. Sadovnick AD. Genetic background of multiple sclerosis. *Autoimmunity reviews* 2012;**11**(3):163-6.

2. Marrie RA, Yu N, Blanchard J, et al. The rising prevalence and changing age distribution of multiple sclerosis in Manitoba. *Neurology* 2010;**74**(6):465-71.

3. D'Hooghe M B, Haentjens P, Nagels G, et al. Alcohol, coffee, fish, smoking and disease progression in multiple sclerosis. *Eur J Neurol* 2012;**19**(4):616-24.

4. Marrie RA, Rudick R, Horwitz R, et al. Vascular comorbidity is associated with more rapid disability progression in multiple sclerosis. *Neurology* 2010;**74**(13):1041-7.

5. Ploughman M, Austin MW, Murdoch M, et al. Factors influencing healthy aging with multiple sclerosis: a qualitative study. *Disabil Rehabil* 2012;**34**(1):26-33.

6. Ploughman M, Austin MW, Murdoch M, et al. The path to self-management: a qualitative study involving older people with multiple sclerosis. *Physiother Can* 2012;**64**(1):6-17.

7. He D, Guo R, Zhang F, et al. Rituximab for relapsing-remitting multiple sclerosis. *Cochrane Database Syst Rev* 2013;**12**:CD009130.

8. Rietberg MB, Brooks D, Uitdehaag BM, et al. Exercise therapy for multiple sclerosis. *Cochrane Database Syst Rev* 2005(1):CD003980.

9. Ploughman M, Austin M, Stefanelli M, et al. Applying cognitive debriefing to pre-test patient-reported outcomes in older people with multiple sclerosis. *Qual Life Res* 2010;**19**(4):483-7.

10. Canada MSo. MS Research Portal. Secondary MS Research Portal 2012. <http://msresearch.ca/> (accessed May 2014).

11. Hobart J, Lamping D, Fitzpatrick R, et al. The Multiple Sclerosis Impact Scale (MSIS-29): a new patient-based outcome measure. *Brain* 2001;**124**(Pt 5):962-73.
12. Hobart JC, Riazi A, Lamping DL, et al. How responsive is the Multiple Sclerosis Impact Scale (MSIS-29)? A comparison with some other self report scales. *J Neurol Neurosurg Psychiatry* 2005;**76**(11):1539-43.
13. Mahoney FI, Barthel DW. Functional Evaluation: The Barthel Index. *Md State Med J* 1965;**14**:61-5.
14. Nicholl L, Hobart J, Dunwoody L, et al. Measuring disability in multiple sclerosis: is the Community Dependency Index an improvement on the Barthel Index? *Mult Scler* 2004;**10**(4):447-50.
15. Cano SJ, O'Connor RJ, Thompson AJ, et al. Exploring disability rating scale responsiveness II: do more response options help? *Neurology* 2006;**67**(11):2056-9.
16. van der Putten JJ, Hobart JC, Freeman JA, et al. Measuring change in disability after inpatient rehabilitation: comparison of the responsiveness of the Barthel index and the Functional Independence Measure. *J Neurol Neurosurg Psychiatry* 1999;**66**(4):480-4.
17. Yeo D, Faleiro, R., Lincoln, N.B. Barthel ADL Index: a comparison of administration methods. *Clin Rehabil* 1995;**9**(1):34-39.
18. Katz PP. Measures of adult general functional status: The barthel index, katz index of activities of daily living, health assessment questionnaire (HAQ), MACTAR patient preference disability questionnaire, and modified health assessment questionnaire (MHAQ). *Arthritis Care and Research* 2003;**49**(5).
19. Marrie RA, Horwitz RI. Emerging effects of comorbidities on multiple sclerosis. *Lancet Neurol* 2010;**9**(8):820-8.

20. Horton M, Rudick RA, Hara-Cleaver C, et al. Validation of a self-report comorbidity questionnaire for multiple sclerosis. *Neuroepidemiology* 2010;**35**(2):83-90.

21. Wade DT, Legh-Smith J, Langton Hewer R. Social activities after stroke: measurement and natural history using the Frenchay Activities Index. *Int Rehabil Med* 1985;**7**(4):176-81.

22. Turnbull JC, Kersten P, Habib M, et al. Validation of the Frenchay Activities Index in a general population aged 16 years and older. *Arch Phys Med Rehabil* 2000;**81**(8):1034-8.

23. Carter J, Mant F, Mant J, et al. Comparison of postal version of the Frenchay Activities Index with interviewer-administered version for use in people with stroke. *Clin Rehabil* 1997;**11**(2):131-8.

24. Einarsson U, Gottberg K, Fredrikson S, et al. Activities of daily living and social activities in people with multiple sclerosis in Stockholm County. *Clin Rehabil* 2006;**20**(6):543-51

25. Ytterberg C, Johansson S, Gottberg K, et al. Perceived needs and satisfaction with care in people with multiple sclerosis: a two-year prospective study. *BMC Neurol* 2008;**8**:36.

26. Godwin M, Streight S, Dyachuk E, et al. Testing the Simple Lifestyle Indicator Questionnaire: Initial psychometric study. *Can Fam Physician* 2008;**54**(1):76-7.

27. Honarmand K, Feinstein A. Validation of the Hospital Anxiety and Depression Scale for use with multiple sclerosis patients. *Mult Scler* 2009;**15**(12):1518-24.

28. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;**67**(6):361-70.

29. Wagnild GM, Young HM. Development and psychometric evaluation of the Resilience Scale. *J Nurs Meas* 1993;**1**(2):165-78.

30. Weinert C. A social support measure: PRQ85. *Nurs Res* 1987;**36**(5):273-7.

- 1
2
3 31. Weinert C, Cudney S, Hill WG. Rural women, technology, and self-management of chronic
4 illness. *Can J Nurs Res* 2008;**40**(3):114-34.
5
6
7
8 32. Long KA, Weinert C. Descriptions and perceptions of health among rural and urban adults
9 with multiple sclerosis. *Res Nurs Health* 1992;**15**(5):335-42.
10
11
12 33. Canada S. Canadian Community Health Survey: Data Liberation Initiative 2010.
13
14 34. Gagne L, Tedds, L., Sullivan, J., Berniaz, K. An introduction to the UT/DLS: Microdata
15 analysis subsetting. Secondary An introduction to the UT/DLS: Microdata analysis
16 subsetting n.d. <http://sda.chass.utoronto.ca/sdaweb/doc/SDAtutorial2008.pdf>.
17
18
19 35. Hawthorne G, Elliott P. Imputing cross-sectional missing data: comparison of common
20 techniques. *Aust N Z J Psychiatry* 2005;**39**(7):583-90.
21
22
23 36. Evans C, Beland SG, Kulaga S, et al. Incidence and prevalence of multiple sclerosis in the
24 Americas: a systematic review. *Neuroepidemiology* 2013;**40**(3):195-210.
25
26
27 37. Canada S. Population by sex and age group Secondary Population by sex and age group
28 2012. <http://www.statcan.gc.ca/tables-tableaux/sum-som/l01/cst01/demo10a-eng.htm>.
29
30
31 38. Buchanan RJ, Chakravorty BJ, Tyry T, et al. Age-related comparisons of people with
32 multiple sclerosis: demographic, disease, and treatment characteristics.
33 *NeuroRehabilitation* 2009;**25**(4):271-8.
34
35
36 39. Giordano A, Ferrari G, Radice D, et al. Self-assessed health status changes in a community
37 cohort of people with multiple sclerosis: 11 years of follow-up. *Eur J Neurol*
38 2013;**20**(4):681-8.
39
40
41 40. Asano M, Duquette P, Andersen R, et al. Exercise barriers and preferences among women
42 and men with multiple sclerosis. *Disabil Rehabil* 2013;**35**(5):353-61.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

41. Kasser SL, Kosma M. Health beliefs and physical activity behavior in adults with multiple sclerosis. *Disabil Health J* 2012;**5**(4):261-8.

42. Jensen MP, Molton IR, Gertz KJ, et al. Physical activity and depression in middle and older-aged adults with multiple sclerosis. *Disabil Health J* 2012;**5**(4):269-76.

43. Krokavcova M, Nagyova I, van Dijk JP, et al. Mastery, functional disability and perceived health status in patients with multiple sclerosis. *Eur J Neurol* 2008;**15**(11):1237-44.

44. Kayes NM, McPherson KM, Schluter P, et al. Exploring the facilitators and barriers to engagement in physical activity for people with multiple sclerosis. *Disabil Rehabil* 2011;**33**(12):1043-53.

45. Chretien JP, Chu LK, Smith TC, et al. Demographic and occupational predictors of early response to a mailed invitation to enroll in a longitudinal health study. *BMC Med Res Methodol* 2007;**7**:6.

46. Manouchehrinia A, Tench CR, Maxted J, et al. Tobacco smoking and disability progression in multiple sclerosis: United Kingdom cohort study. *Brain* 2013;**136**(Pt 7):2298-304.

47. D'Hooghe M B, Nagels G, De Keyser J, et al. Self-reported health promotion and disability progression in multiple sclerosis. *J Neurol Sci* 2013;**325**(1-2):120-6.

48. Marrie RA, Horwitz RI, Cutter G, et al. Smokers with multiple sclerosis are more likely to report comorbid autoimmune diseases. *Neuroepidemiology* 2011;**36**(2):85-90.

49. Weiland TJ, Hadgkiss EJ, Jelinek GA, et al. The association of alcohol consumption and smoking with quality of life, disability and disease activity in an international sample of people with multiple sclerosis. *J Neurol Sci* 2014;**336**(1-2):211-9.

Figure Legends:

Figure 1: Survey Respondents

Figure 2: Number of respondents mapped by postal code

Figure 3: Frequency of Barthel Index score categories: A score of 100 on the Barthel Index suggests that an individual is independently ambulatory, can ascend and descend stairs on their own, is continent and able to perform ADLs without help. Fifteen percent of this cohort were independent. A Barthel Index score of 91-99 suggests slight dependence (16%); 61-90 suggests moderate dependence (47%); 21-60 suggests severe dependence (16%) and 0-20 represents complete dependence (7%).

Figure 4: Frequency of Frenchay Activities Index score categories

The Canadian Survey of Health, Lifestyle and Aging with Multiple Sclerosis; Methodology and Initial Results

Michelle Ploughman PT, PhD*, Serge Beaulieu**, Chelsea Harris BKin MPH*, Stephen Hogan BKin*, Olivia J. Manning*, Penelope W. Alderdice PhD*, John D. Fisk PhD[§], A. Dossa Sadovnick PhD[†], Paul O'Connor, MD, FRCPC[‡], Sarah A. Morrow, MD[†], FRCPC, Luanne M. Metz MD, FRCPC[#], Penelope Smyth MD, FRCPC[¶], Nancy Mayo, PhD^{§§}, Ruth Ann Marrie, MD, PhD^{††}, Katherine B. Knox, MD, FRCPC^{‡‡}, Mark Stefanelli MD FRCPC^{¶¶}, Dr. Marshall Godwin, MD^{##}

*Recovery and Performance Laboratory, Rehabilitation Research Unit, Faculty of Medicine, Memorial University, St. John's NL Canada

**Eastern Health Authority, St. John's NL Canada

[§]Dalhousie University, Halifax NS, Canada

[†]Department of Medical Genetics and Division of Neurology, Faculty of Medicine University of British Columbia, Vancouver BC, Canada

[‡]Department of Neurology, St. Michaels Regional Hospital, Toronto ON, Canada

[†]London Health Science Centre, London ON, Canada

[¶]Department of Neurology, University of Alberta, Edmonton AB, Canada

[¶]Clinical Neurosciences, University of Calgary, Calgary AB, Canada

^{§§}Clinical Epidemiology, McGill University, Montreal QC, Canada

^{††}Departments of Internal Medicine and Community Health Sciences, University of Manitoba, Winnipeg MB, Canada

^{‡‡}Department of Physical Medicine and Rehabilitation, College of Medicine, University of Saskatchewan, Saskatoon SK, Canada

^{¶¶}Neurology, Faculty of Medicine, Memorial University, St. John's NL Canada

^{##}Primary Health Care Research Unit, Faculty of Medicine, Memorial University, St. John's NL, Canada

Corresponding Author/Research Site

Dr. Michelle Ploughman, Recovery and Performance Laboratory, Rehabilitation Research Unit, Faculty of Medicine, Memorial University, Rm 400, L.A. Miller Centre, 100 Forest Rd., St. John's NL Canada. Ph 709 777 2099; fax 709 777 8702; email mploughm@mun.ca

Key Words: aging; diet; exercise; health behaviours; multiple sclerosis; neurodegenerative disease; older persons; quality of life

Short Title: Health Lifestyle and Aging with Multiple Sclerosis

Tables: 3

Figures: 4

For peer review only

ABSTRACT

Objective

People with multiple sclerosis (MS) are living longer so strategies to enhance long term health and quality of life are garnering more interest. We aimed to create a profile of aging with MS in Canada by recruiting 1250 (5% of the Canadian population over 55 with MS) participants and focusing data collection on health and lifestyle factors, disability, participation and quality of life in order to determine factors associated with healthy aging.

Design

National multi-centre postal survey

Setting

Recruitment from Canadian MS clinics, MS Society of Canada chapters, and newspaper advertisements

Participants

People aged 55 years or older with MS symptoms more than 20 years

Outcome Measures

Validated outcome measures and custom-designed questions examining MS disease characteristics, living situation, disability, co-morbid conditions, fatigue, health behaviours, mental health, social support, impact of MS, and others.

Results

Of the 921 surveys, 743 were returned (80.7% response rate). Participants (mean age 64.6 ±6.2 years) reported living with MS symptoms for an average of 32.9±9.5 years and 28.6% were either wheelchair users or bedridden. There was only 5.4% missing data and 709 respondents provided optional qualitative information. According to data derived from the 2012 Canadian Community Health Survey of Canadians over 55 years of age, older people with MS from this survey sample are about 8 times less likely to be employed full-time. Older people with MS were less likely to engage in regular physical activity (26.7%) compared to typical older Canadians (45.2%). However, they were more likely to abstain from alcohol and smoking.

Conclusion

Despite barriers to participation, we were able to recruit and gather detailed responses (with good data quality) from a large proportion of older Canadians with MS. The data suggests that this sample of older people with MS are less likely to be employed, are less active and more disabled than other older Canadians.

Strengths and limitations of this study

- This unique cohort provides new insights on aging with MS. With an average age about 65 years, our respondents are 10 to 25 years older than cohorts examined in previous studies of older people with MS.
- Our recruitment methods and a very high survey return rate allowed us to obtain data from about 3% of our target population.
- With only about 10% of our sample being employed in any capacity, our results point to a need to understand retirement and employment in aging with MS.
- The cross sectional design limits our ability to assess change and the effects of variables on individual outcomes over time.
- The low sampling of persons in nursing homes and assisted living situations highlights the need for specialized recruitment strategies to study this sub-set of people living with MS.

This research was supported by a Canadian Institutes of Health Research (CIHR) Post-doctoral Fellowship, a Newfoundland and Labrador Centre for Applied Health Research, Healthy Aging Research Program Project Grant, the Health Care Foundation (Eastern Health Authority) Research Project Grant, and the Physiotherapy Foundation of Canada Alberta Research Award.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

Multiple Sclerosis (MS) is a highly unpredictable and heterogeneous autoimmune disease diagnosed between the ages of 18-40 years. Affecting women more than men, the cause of MS is unknown, however research suggests that a combination of environmental factors and genetic predisposition may be the trigger for disease onset [1]. The number of people with MS over the age of 55 is increasing, possibly due to improved longevity and more effective treatments [2]. On the other hand, several authors have shown that smoking [3] and cardiovascular disease [4] accelerate disease progression. People with MS are interested in strategies that minimize disability and enhance role participation and quality of life [5]. The uncertainty surrounding MS progression along with aging MS patients warrants the need for investigation into potential disease modifying health behaviours. Older individuals who have lived with MS for many years may provide insight into factors that contribute to living a long and healthy life with MS [6].

Older people with MS are often excluded from MS research. Studies involving clinical drug trials [7] and exercise interventions [8], in particular, target patients with mild relapsing-remitting disease. Furthermore, because they may not require intensive medical oversight associated with taking disease-modifying drugs, older patients may not attend MS clinics as often as those recently diagnosed. The older MS cohort are also more likely to be disabled and living in an institutional setting, making recruitment challenging. In addition, disability-related impairments such as those affecting vision, cognition and coordination complicate survey and patient-reported outcome administration [9].

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

In order to create a profile of aging with MS in Canada, we developed a national cross-sectional survey, based on previous qualitative research [5 6], using validated patient-reported outcome measures [9]. We focused on health and lifestyle factors, disability, participation and quality of life, targeting people 55 years of age and older with MS symptoms for 20 or more years. By partnering with major MS clinic registries and MS society chapters, we hypothesized that we would be able to reach a representative sample of about 5% of the ‘aging with MS in Canada’ cohort.

METHODS

Participants and Sampling: We targeted Canadians, aged 55 years or older, who had had self-reported MS symptoms for 20 or more years. Prior to recruitment, the study was approved by 11 health research ethics boards in eight Canadian provinces. Between May 2011 and December 2012, participants were recruited by telephone from lists of eligible MS clinic patients and by public advertising through the MS Society of Canada Research Internet Portal [10], local MS Society chapters and their newsletters, and newspaper advertisements. A confirmed diagnosis from a neurologist was available for participants recruited through the MS clinic databases; self-identified participants provided a diagnosis date and neurologist name to confirm their diagnosis. Rather than calculating a sample size, we used an exhaustive sampling technique to obtain a large and representative sample; for example repeatedly targeting regional newspapers, following-up at study sites, making formal and informal contacts with MS Society Chapter support groups.

After obtaining consent using a standard script, we mailed participants the questionnaire, with an information sheet, a toll-free telephone helpline number and email address and a prepaid

reply envelope. We requested that participants return the completed survey in 2 weeks; non-responders were telephoned for follow-up at 4 weeks. Participants who wished to complete the survey by telephone were encouraged to do so. Each questionnaire was coded to maintain anonymity.

Survey Design: The questionnaire consisted of validated patient-reported outcome tools and additional custom-designed questions chosen based on pilot research in a small sample of older people with MS [9]. We determined during this step that participants preferred a paper rather than online version of the survey as well as larger 14-point font, divided into sections that grouped similar questions together. Respondents were instructed to complete the survey in these sections (chapters) taking a break in between. Since we previously observed confusion with questions involving temporal parameters, questions regarding activity involvement that followed the same timeline (e.g. weekly, monthly, or yearly) were placed in sequence. To further examine the acceptability of the survey and to minimize skipped or misunderstood items, we asked the first 14 participants (11 returns) to provide feedback on the content, format and response burden. This feedback was integrated into the final version of the questionnaire but the responses were not included in the final data.

In order to include people with MS in the province of Quebec, Canada, the entire survey including the cover letter was translated into french by one independent professional translator. We also obtained french versions of previously validated tools. Four native french speakers involved in MS research made corrections to the initial translation in sequence. The corrections were then consolidated by a fifth french speaking MS researcher. The advertisement and

consent scripts were translated by a native french speaking MS researcher and reviewed and corrected by a second.

Survey Content: The final survey consisted of five chapters (**Table 1**) ending with open response questions for participants to provide additional comments. Where appropriate, we obtained permission from the developers to use previously validated tools.

We used the Multiple Sclerosis Impact Scale (MSIS-29), the Barthel Index and the Co-morbidity Questionnaire to assess health. The MSIS-29 consists of 29 questions: 20 address the physical impact and nine address the psychological impact of MS generating separate or combined scores as a percentage of 100. Higher scores indicate greater impact of MS [11]. The MSIS-29 physical component correlates with the Barthel Index (0.71) and the psychological component with EQ-5D (0.68). It is reported to have good responsiveness with large to moderate effect sizes [12]. The Barthel Index measures function and mobility in activities of daily living (ADL) in ten areas: feeding, bathing, grooming, dressing, bowel control, bladder control, toileting, transfers, ambulation and stair climbing [13]. Items are rated in terms of whether individuals can perform activities independently, with some assistance, or are completely dependent, with a total score ranging from 0 (complete assistance required) to 100 (independent). The Barthel Index has been tested in people with MS [14-16] and is reliable whether administered by telephone, in-person or by mail [17] with effect sizes comparable to the Functional Independence Measure (0.34) [18]. The list of co-morbidities used in this survey was adapted from Marrie and Horwitz [19 20] in consultation with Dr. Marrie (personal communication).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

We determined the degree of participation in instrumental activities of daily living (iADL) using the Frenchay Activities Index (FAI) and individual components of lifestyle (exercise, diet, smoking, alcohol, and stress) using the Simple Lifestyle Indicator Questionnaire (SLIQ). The FAI assesses the frequency of participation in household chores, social activities, work and leisure for a summed raw score of 15 to 60; higher score indicating more frequent participation [21]. The FAI has excellent test-retest reliability as a mailed questionnaire (0.96) [22] and correlates well with face-to-face interview scores [23]. The FAI has been mainly used in stroke but has also been used in large MS population studies [24 25]. The SLIQ was developed to measure the individual components of lifestyle that affect cardiovascular disease [26]. Initial content and face validity were assessed by the developers among 136 people attending a family practice clinic[26] and SLIQ scores were highly correlated with those of blinded reviewers (0.77). We were also interested in mental health, coping and social support in aging with MS. The Hospital Anxiety and Depression Scale (HADS) is a self-assessment scale found to reliably screen for depression and anxiety in multiple sclerosis patients [27]. It is divided into an Anxiety subscale (HADS-A) and a Depression subscale (HADS-D) each containing seven intermingled items scored out of 21; higher score indicating greater symptoms [28]. A threshold score of 8 or higher on HADS subscales had high sensitivity (Depression 90%, Anxiety 88.5%) and specificity (Depression 87.3%, Anxiety 80.7%) compared to structured interview using DSM-1 criteria among people with MS [27]. Resilience, the ability to overcome adversity, was measured using the 25 item Resilience Scale. Responses to the test questions range from 1 (disagree) to 7 (agree) and the summed score can be further interpreted as, low resilience (<120), moderate resilience (125-145), and high resilience (>145) [29] and has been validated in older community-

living individuals with high internal consistency reliability (0.91)[29] . The Personal Resource Questionnaire (PRQ-2000) measures social support and consists of 15 statements with which the subject agrees or disagrees based on a seven-point Likert scale; higher scores indicating higher support (e.g. I belong to a group in which I feel important 1-Strongly Disagree to 7-Strongly Agree). PRQ-2000 demonstrates good internal consistency (0.87 to 0.93), adequate construct validity (moderate correlations with anxiety -0.37, neuroticism -0.28 and extroversion 0.32) and good predictive and construct validity in multiple chronic disease data sets [30]. Mailed versions have been used in community-living women with chronic health conditions [31] and adults with MS [32].

Table 1: Survey Content

CHAPTER	DOMAINS	MEASUREMENT TOOLS
You and Your MS (5 pages)	Demographics	Postal code, gender, height, weight, age, finances and education
	MS Disease Characteristics	Symptom onset, time since diagnosis, and type of MS
	Environment	Living situation, accessibility and home modifications
Your Health (6 pages)	Health-Related Quality of Life	Visual analogue scale of perceived health status [9] Multiple Sclerosis Impact Scale-29 [11]
	Physical Health and Disability	Barthel Index [12] Co-Morbidity Questionnaire [17]
	Health Care Services	Use and satisfaction with health care services, medications and complementary/alternative health.
Your Activities (7 pages)	Participation and instrumental Activities of Daily Living	Frenchay Activities index [19]; Employment and impact of fatigue
	Lifestyle	Simple Lifestyle Indicator Questionnaire (diet, exercise, stress, smoking, alcohol) [24]
Your Thoughts and Feelings (8 pages)	Mental/ Cognitive Health	Hospital Anxiety and Depression Scale (HADS) Resilience Scale [25] Problems with concentration and memory
Your Social Life (4 pages)	Social Support	Personal Resource Questionnaire-2000 [30]

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Canadian comparison data: Comparative data was collected from the Canadian Community Health Survey (CCHS) 2012 version; a national cross-sectional survey that gathers information about health, social and economic status and lifestyle behaviours of the population. Data was extracted from the CCHS database using the University of Toronto Data Library Service (DLS) [33]. A CCHS sample was selected based on age (>55 years). Data was collected from CCHS questions that matched question from our survey (e.g. mobility from the Barthel Index and lifestyle from the SLIQ). Outputs (ie. highest level of education, employment status) were analyzed using Survey Documentation & Analysis (SDA) software [34]. Frequency tables were created for variables of interest.

Data Cleaning and Imputation: We custom designed the database application to house project data and minimize error during data imputation. Missing data was imputed using three techniques: (1) deductive imputation, (2) person mean substitution, and (3) ‘hot deck’ imputation [35]. The deductive technique was used to impute missing data when there was sufficient comments and answers to related questions. If less than half of the data was missing on a section, scale, or subscale, person mean substitution (PMS) was used. PMS takes the mean score of the particular participant on that section, scale, or subscale and uses the mean as the value for the missing data. If greater than half of the data was missing on a section, scale, or subscale, ‘hot deck’ imputation was used [35]. This involved taking a value from the "hot deck" or in other terms, finding another participant of the same gender, age (± 5 years), education level, and living in the same province. Hot deck imputation was selected for this study because it is proven to be the most efficient method of data imputation as it is more robust in large samples when

1
2
3 compared to other imputation methods [35]. The exact scores for each question from the
4
5 substitute participant were imputed for the missing values for the participant of interest.
6
7
8 Variables that could not be imputed and remained empty included, from **Table 1**, Demographics
9
10 (e.g. weight, height, finances), MS Disease Characteristics, Environment, and Health Care
11
12 Services.
13

14
15
16
17 **Data Analysis:** Data was analyzed using IBM SPSS software v20. Frequencies and descriptive
18
19 statistics (mean, SD and data range) were used to determine subject characteristics.
20
21
22
23
24
25
26

27 RESULTS

28 *Survey Response*

29
30
31 We aimed to recruit 1250 participants for the study and through exhaustive sampling we
32
33 contacted 921 (74% of target) people with MS who met inclusion criteria. Seven hundred and
34
35 forty three participants returned the survey (**Figure 1**; 80.7% response rate). Survey response by
36
37 province is illustrated in **Figure 2**. Of the respondents, 90.4% (n=672) were identified through
38
39 provincial MS clinics while the remaining 71 self-identified by responding to newspaper ads and
40
41 word of mouth. Forty-five people requested help to complete the survey, and 11 participants
42
43 asked that the entire survey be completed by phone due to writing or vision difficulties or both.
44
45 Additional content was provided by 709 participants in the open-ended questions at the end of
46
47 the survey.
48
49
50
51
52

53 Only 563 of 104,763 cells contained missing data (5.37%). Almost every survey
54
55 contained one or two random missing items; in some cases answers to these could be inferred
56
57
58
59
60

using deductive reasoning (67/104,763=0.06%). We observed no pattern of missing data other than body weight (which we did not impute). The PMS technique was required for 170 cells (170/94,763=0.16%) and the ‘hot deck’ technique was required in 14 surveys (324 cells/104,763=0.31%). Cases of consecutive missing data seemed to be mainly due to pages adhering together resulting in questions being overlooked.

Insert Figure 1 about here

Insert Figure 2 about here

Sample Characteristics

Participants ranged in age from 55 to 88 years with a mean age of 64.6 years (±6.2). They described experiencing MS symptoms for 20 to 73 years with a mean duration of 32.9 years (±9.5). The respondents reported that the average time from onset of symptoms to diagnosis by a neurologist was 8.1 years (± 9.4) with a range from 0 to 47 years. The number of women outnumbered men 3.5:1 (Table 2).

Sixty percent reported at least one year of post-secondary education, and 90% were either retired or unemployed. About twenty-eight percent of the sample (n=212) reported that they were no longer able to walk and either used a wheelchair or were bedridden. The characteristics of the sample are fully described in Table 2.

Respondents were categorized by level of disability based on results of the Barthel Index (Figure 3) [18]. We measured activity and participation using the FAI which is divided into household (domestic) activity, leisure activity and outdoor activity (Figure 4). Nearly half

(50.6%) of the sample frequently participated in household activities but only 13.5% and 27.9% reported that they frequently engaged in leisure activities and outdoor activities respectively.

Table 2: Sample Characteristics

Characteristic		N (%)
Years with MS Symptoms	20-29	324 (43.6)
	30-39	245 (33.0)
	40-49	135 (18.2)
	50-59	31 (4.2)
	60 or more	7 (0.9)
Initial MS Diagnosis	Relapsing-Remitting	386 (52.0)
	Primary Progressive	99 (13.3)
	Secondary Progressive	65 (8.7)
	Progressive-Relapsing	16 (2.2)
	Benign	43 (5.8)
	Unknown	126 (17.0)
Living situation	Living in a private home	715 (96.2)
	Living in an assisted living complex	16 (2.2)
	Living in long term care	11 (1.5)
	Home Care	146 (19.7)
	Equipment	200 (26.9)
	Home Modifications	73 (9.8)
Private Insurance (Health/Disability)	Yes	393 (52.9)
	No	350 (47.1)

Insert Figure 3 about here

Insert Figure 4 about here

Older people with MS compared to typical older Canadians

A comparison sample of older people (>55) in Canada was derived from the 2012 Canadian Community Health Survey (CCHS; **Table 3**). CCHS respondents over the age of 55 years primarily lived with their spouse or partner (52.5%), with their spouse or partner and children (11.3%), or alone (22.7%). Only approximately one percent of this sample were living

with their parents. According to data derived from the CCHS, older people with MS from our survey sample have less post-secondary education and are about 8 times less likely to be employed full-time. About half of the MS survey sample have major mobility challenges (unable to walk outdoors, use a wheelchair or are bedridden) compared to only 9% in the CCHS sample. Older people with MS were more likely to engage in physical activity (69.4%) compared to typical older Canadians (45.3%). They were also more likely to abstain from alcohol and smoking.

Table 3: Comparison of Sample Characteristics to Others Aging in Canada

Characteristic		Study sample (%)	CCHS sample (%)
Age	55-64	414 (55.7)	4,671,364.6 (48.9)
	65-74	274 (36.9)	2,891,671.1 (30.3)
	75+	55 (7.4)	1,989,358.6 (20.8)
Gender	Males	166 (22.3)	4,568,522.0 (47.8)
	Females	577 (77.7)	4,983,872.3 (52.2)
Education	Less than high school	74 (10.0)	1,241,946 (15.2)
	High School	222 (29.9)	1,083,618 (13.3)
	1-2 year post-secondary	189 (25.4)	4,088,252 (4.9)
	Complete Post-graduate	258 (34.7)	5,442,586 (66.6)
Employment Status	Retired	504 (67.8)	2,878,200 (46.3)
	Unemployed	166 (22.3)	302,942 (4.8)
	Employed Full-time	30 (4.0)	2,068,571 (33.3)
	Employed Part-time	19 (2.6)	698,125 (11.2)
	Employed Casual	24 (3.2)	266,419 (4.3)
Mobility	Walk independently with or without cane outdoors	351 (47.2)	8,004,425 (90.9)
	Walk with cane or help indoors	180 (24.2)	593,584 (6.7)
	Wheelchair	161 (21.7)	
	Unable to move	51 (6.9)	204,744 (2.3)
Lifestyle	Moderate to Vigorous Exercise	193 (25.9)	1,764,305 (20.7)
	Some physical activity	316 (42.5)	2,083,695 (24.5)
	No physical activity	234 (31.5)	4,652,856 (54.7)
	No alcohol use	385 (51.8)	2,226,486 (26.3)
	14 or less drinks per week	344 (46.3)	1,242,238 (14.6)
	>14 drinks per week	14 (1.9)	5,009,245 (59)
	Non-Smokers	676 (91.0)	7,422,711 (84.4)
	Never smoked	272 (36.6)	
	Quit	404 (54.4)	
	Smokers	67 (9.0)	1,372,885 (15.6)

DISCUSSION

The purpose of this national cross-sectional survey was to describe aging with MS in Canada that would provide the data necessary to determine, in future analysis, the influence of personal, health and environmental factors on quality of life and disability among older people with MS. Incidence and prevalence estimates of MS in Canada are amongst the highest worldwide [36] and a shift in peak age prevalence of MS to older ages suggesting that persons with MS in Canada are living longer [2]. Interventions to promote successful aging with MS may take on more importance over the next decade since quality-of-life and productivity in advanced age is a desired outcome for most [6]. To our knowledge this is the most comprehensive study focusing on examination of health and lifestyle factors specifically among older persons with MS.

The prevalence of MS in Canada has been estimated to be 240 per 100,000 [36]. The CCHS estimates that there are 93,535 Canadians living with MS [33] and since Statistics Canada estimates that people over 55 make up about 27.7% of the population [37], there could be approximately 25,000 people with MS over 55 years of age. We attempted to contact as many older people with MS as possible and our sample (n=743) likely represents about 3% of this group, less than our initial 5% target. Data quality was very good with only 5.4% missing data and 709 respondents providing optional qualitative data. The high rates of response and completion and may be due to pre-screening of potential respondents at clinic sites. Only 7.5% of respondents required assistance by the research team but we were unable to discern survey completion by proxy.

With an average age about 65 years, our respondents are 10 to 25 years older than cohorts examined in previous studies of older people with MS [38-44]. Compared to these previous

1
2
3 studies, which reported the number of years since MS diagnosis ranging from 8-19 years [39 40],
4
5 our participants have been diagnosed with MS for 24.8 (± 10.0) years. Our respondents' level of
6
7 education aligns with most other studies of people with MS [39-41 43] .The group is fairly well-
8
9 educated which is not surprising considering characteristics of people who respond to surveys in
10
11 general [45]. When compared to the Canadian population [33], fewer of our respondents had
12
13 advanced education including post-secondary degrees and post-graduate degrees. Furthermore,
14
15 only about 10% of our sample was employed in any capacity which is substantially lower than
16
17 other studies of older people with MS [38 40 43]. Since maintaining employment is associated
18
19 with better quality-of-life, at least in younger people with MS [43], factors associated with
20
21 employment and retirement as one ages with MS will be an important area of future study.
22
23
24
25
26

27 More than a quarter of our cohort were severely disabled (28.6%); over ten times that of
28
29 aging counterparts in Canada (2.3%). They either were restricted to ambulation with a
30
31 wheelchair or walking with physical assistance indoors or were bedridden; an interesting finding
32
33 given that most respondents lived in private residences (96.2%) without home care, equipment or
34
35 home modifications. Since only about half of respondents had private health insurance, it is
36
37 possible that most people manage at home with few resources outside of the publicly-funded
38
39 health system and the care provided by spouses and children. It was interesting that only a small
40
41 percentage of survey respondents engaged in leisure or outdoor activity (Figure 4). Since the
42
43 proportion of respondents living in assisted-living and long-term care in this study was small, our
44
45 sample may not be representative of older people with MS living in these settings.
46
47
48
49

50 In terms of lifestyle behaviours, the participants in this study were more physically active
51
52 and less likely to drink alcohol or smoke compared to their Canadian counterparts [33]. In large
53
54 cohorts of people with MS in the United Kingdom (n=895) and Belgium (n=1372), healthy
55
56
57
58
59
60

lifestyle behaviours are associated with less severe and less rapid disability progression [46 47]. Poor lifestyle behaviors, specifically smoking, have been associated with increased risk of comorbid diseases [48] and significantly lower HRQOL when compared to people with MS who have never smoked [49]. Future studies will examine the role of health behaviors and disability in this group of older people with MS.

Although this unique cohort may provide new insights on aging with MS, there are some study limitations. The cross sectional design limits our ability to assess change and the effects of variables on individual outcomes over time. By nature of the volunteer survey design, our sample may be biased in that active participants were more likely to respond. Even though we provided assistance over the phone, those with significant disability who required assistance may have declined. The low sampling of persons in nursing homes and assisted living situations and those older than 75 years highlights the need for specialized recruitment strategies to study this sub-set of people living with MS. Although preliminary testing indicated that participants preferred paper, a web-based version may have improved our recruitment. We also have compared our sample to the Canadian Census CCHS sample. The survey questions do not perfectly align so the values are estimates only. A final limitation of the survey was that we selected the Simple Lifestyle Indicator Questionnaire to assess health behaviours because of its short, simple design, so the data collected lacks detail on diet behaviours which will limit future analysis.

Conclusion

Despite barriers to participation, we were able to recruit and gather detailed responses (with good data quality) from a large proportion of older Canadians with MS. Even with extensive pretesting, national collaboration, exhaustive sampling and high return rates, our

recruitment (n=743) was less than our target (n=1250). However, we were able to recruit a large proportion of people who were severely disabled even though only 7.5% of respondents requested assistance. The data suggests that this sample of older people with MS are less likely to be employed, are more disabled but have more positive health behaviours than other older Canadians.

Acknowledgements

Support for this project was received from the MS Society of Canada and local chapters; provincial MS clinic neurologists, MS nurses, administrative and research staff; and the Rehabilitation and Continuing Care Program, Eastern Health Authority. We specifically acknowledge the work of site coordinators: Trudy Campbell, Heather Rosehart, Amy Cuthbertson, Darren Nickel, Pam Dumont, Madonna de Lemos, Deborah Goldberg, Irene Yee, Nicholas Hall, Maureen Perera, Sarah Williamson, Carolina Moriello and Vanessa Bouchard.

Contributorship Statement

Concept development and funding applications- MP, JDF, MG
Ethics applications and participant recruitment-MP, OJM, CH, SH, JDF, ADS, PO, SAM, LMM, PS, NM, RAM, MS
Data collection, management and analysis-MP, OJM, SB, CH, SH, PWA
Manuscript preparation and /or review-MP, OJM, CH, JDF, ADS, PO, SAM, LMM, PS, NM, RAM, MS, MG

Competing Interests

Dr. Michelle Ploughman was supported by a Canadian Institutes for Health Research (Institute of Aging) Postdoctoral Fellowship. Her research was supported by the Newfoundland and Labrador Centre for Applied Health Research, Health Care Foundation, and Physiotherapy Foundation of Canada.

Serge Beaulieu, Stephen Hogan, Dr. Penelope Alderdice, Dr. A. Dessa Sadovnick, Dr. Luanne M. Metz, and Dr. Nancy Mayo have no disclosures or conflicts of interest to report.

Chelsea Harris and Olivia Manning were supported by endMS Research and Training Network summer studentships.

Dr. John Fisk is the Director of the endMS Atlantic Regional Research and Training Centre, which is funded by the Multiple Sclerosis Society of Canada. He receives research funding from the Canadian Institutes of Health Research (CIHR) and in the past has received grants, honoraria and consultation fees from Astra- Zeneca, Bayer, Biogen-Idec Canada, Heron Evidence Development Limited, Hoffmann-La Roche, MAPI Research Trust, Novartis, Sanofi-Aventis, Serono Canada, and QualityMetric Inc.

Dr. Paul O'Connor has received consulting fees and/or research support for MS trials from Actelion, Bayer, Biogen Idec, BioMS, Cognosci, Daiichi Sankyo, EMD Serono, Genentech, Genmab, Novartis, Roche, Sanofi Aventis, Teva and Warburg Pincus.

Dr. Sarah Morrow has received research support from Biogen Idec and Novartis Canada, and received support for speaking engagement and advisory panels for Bayer, Biogen Idec, EMDSerono and Teva Neurosciences and Novartis Canada.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Dr. Penelope Smyth has served on scientific advisory boards for Novartis and EMD Serono. She has received funding for travel to pharmaceutically sponsored events by EMD Serono. She has received honoraria for speaking, facilitating and chairing from Novartis and EMD Serono as well as from the MS Society of Canada. She has received academic research support for medical education research from the Arnold P. Gold Foundation, and the Teaching and Learning Enhancement Fund from the University of Alberta.

Dr. Ruth Ann Marrie receives research funding from: Canadian Institutes of Health Research, Multiple Sclerosis Society of Canada, Multiple Sclerosis Scientific Research Foundation, Rx & D Health Research Foundation, and has conducted clinical trials funded by Sanofi-Aventis.

Dr. Katherine Knox received grant funding from Saskatoon City Hospital Foundation and MS Society of Canada, educational CME event grant funding from Bayer HealthCare Pharmaceuticals, honoraria /travel grant from Biogen and Teva. She was a member of the CIHR and MS Society of Canada Scientific Expert Working Group on CCSVI and MS.

Dr. Mark Stefanelli has served on advisory boards for Biogen Idec and Novartis and has been provided speaker/travel support by Teva Neuroscience. His research was supported by Genzyme and Biogen.

Dr. Marshall Godwin's research is funded by the Canadian Institutes of Health Research, Public Health Agency of Canada, Health Canada, and Newfoundland and Labrador Centre for Applied Health Research. He reports no conflicts of interest.

Funding

This research was supported by a Canadian Institutes of Health Research (CIHR) Post-doctoral Fellowship, a Newfoundland and Labrador Centre for Applied Health Research, Healthy Aging Research Program Project Grant, the Health Care Foundation (Eastern Health Authority) Research Project Grant, and the Physiotherapy Foundation of Canada Alberta Research Award.

Data Sharing

Additional data can be obtained by contacting Dr. Michelle Ploughman at Michelle.Ploughman@med.mun.ca.

References

1. Sadovnick AD. Genetic background of multiple sclerosis. Autoimmunity reviews 2012;**11**(3):163-6.

2. Marrie RA, Yu N, Blanchard J, et al. The rising prevalence and changing age distribution of multiple sclerosis in Manitoba. Neurology 2010;**74**(6):465-71.

3. D'Hooghe M B, Haentjens P, Nagels G, et al. Alcohol, coffee, fish, smoking and disease progression in multiple sclerosis. Eur J Neurol 2012;**19**(4):616-24.

4. Marrie RA, Rudick R, Horwitz R, et al. Vascular comorbidity is associated with more rapid disability progression in multiple sclerosis. Neurology 2010;**74**(13):1041-7.

5. Ploughman M, Austin MW, Murdoch M, et al. Factors influencing healthy aging with multiple sclerosis: a qualitative study. Disabil Rehabil 2012;**34**(1):26-33.

6. Ploughman M, Austin MW, Murdoch M, et al. The path to self-management: a qualitative study involving older people with multiple sclerosis. Physiother Can 2012;**64**(1):6-17.

7. He D, Guo R, Zhang F, et al. Rituximab for relapsing-remitting multiple sclerosis. Cochrane Database Syst Rev 2013;**12**:CD009130.

8. Rietberg MB, Brooks D, Uitdehaag BM, et al. Exercise therapy for multiple sclerosis. Cochrane Database Syst Rev 2005(1):CD003980.

9. Ploughman M, Austin M, Stefanelli M, et al. Applying cognitive debriefing to pre-test patient-reported outcomes in older people with multiple sclerosis. Qual Life Res 2010;**19**(4):483-7.

10. Canada MSo. MS Research Portal. Secondary MS Research Portal 2012. <http://msresearch.ca/> (accessed May 2014).

- 1
2
3 11. Hobart J, Lamping D, Fitzpatrick R, et al. The Multiple Sclerosis Impact Scale (MSIS-29): a
4
5 new patient-based outcome measure. *Brain* 2001;**124**(Pt 5):962-73.
6
7
- 8 12. Hobart JC, Riazi A, Lamping DL, et al. How responsive is the Multiple Sclerosis Impact
9
10 Scale (MSIS-29)? A comparison with some other self report scales. *J Neurol Neurosurg*
11
12 *Psychiatry* 2005;**76**(11):1539-43.
13
14
- 15 13. Mahoney FI, Barthel DW. Functional Evaluation: The Barthel Index. *Md State Med J*
16
17 1965;**14**:61-5.
18
19
- 20 14. Nicholl L, Hobart J, Dunwoody L, et al. Measuring disability in multiple sclerosis: is the
21
22 Community Dependency Index an improvement on the Barthel Index? *Mult Scler*
23
24 2004;**10**(4):447-50.
25
26
- 27 15. Cano SJ, O'Connor RJ, Thompson AJ, et al. Exploring disability rating scale responsiveness
28
29 II: do more response options help? *Neurology* 2006;**67**(11):2056-9.
30
31
32
- 33 16. van der Putten JJ, Hobart JC, Freeman JA, et al. Measuring change in disability after
34
35 inpatient rehabilitation: comparison of the responsiveness of the Barthel index and the
36
37 Functional Independence Measure. *J Neurol Neurosurg Psychiatry* 1999;**66**(4):480-4.
38
39
- 40 17. Yeo D, Faleiro, R., Lincoln, N.B. Barthel ADL Index: a comparison of administration
41
42 methods. *Clin Rehabil* 1995;**9**(1):34-39.
43
44
- 45 18. Katz PP. Measures of adult general functional status: The barthel index, katz index of
46
47 activities of daily living, health assessment questionnaire (HAQ), MACTAR patient
48
49 preference disability questionnaire, and modified health assessment questionnaire
50
51 (MHAQ). *Arthritis Care and Research* 2003;**49**(5).
52
53
- 54 19. Marrie RA, Horwitz RI. Emerging effects of comorbidities on multiple sclerosis. *Lancet*
55
56 *Neurol* 2010;**9**(8):820-8.
57
58
59
60

20. Horton M, Rudick RA, Hara-Cleaver C, et al. Validation of a self-report comorbidity questionnaire for multiple sclerosis. *Neuroepidemiology* 2010;**35**(2):83-90.

21. Wade DT, Legh-Smith J, Langton Hewer R. Social activities after stroke: measurement and natural history using the Frenchay Activities Index. *Int Rehabil Med* 1985;**7**(4):176-81.

22. Turnbull JC, Kersten P, Habib M, et al. Validation of the Frenchay Activities Index in a general population aged 16 years and older. *Arch Phys Med Rehabil* 2000;**81**(8):1034-8.

23. Carter J, Mant F, Mant J, et al. Comparison of postal version of the Frenchay Activities Index with interviewer-administered version for use in people with stroke. *Clin Rehabil* 1997;**11**(2):131-8.

24. Einarsson U, Gottberg K, Fredrikson S, et al. Activities of daily living and social activities in people with multiple sclerosis in Stockholm County. *Clin Rehabil* 2006;**20**(6):543-51

25. Ytterberg C, Johansson S, Gottberg K, et al. Perceived needs and satisfaction with care in people with multiple sclerosis: a two-year prospective study. *BMC Neurol* 2008;**8**:36.

26. Godwin M, Streight S, Dyachuk E, et al. Testing the Simple Lifestyle Indicator Questionnaire: Initial psychometric study. *Can Fam Physician* 2008;**54**(1):76-7.

27. Honarmand K, Feinstein A. Validation of the Hospital Anxiety and Depression Scale for use with multiple sclerosis patients. *Mult Scler* 2009;**15**(12):1518-24.

28. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;**67**(6):361-70.

29. Wagnild GM, Young HM. Development and psychometric evaluation of the Resilience Scale. *J Nurs Meas* 1993;**1**(2):165-78.

30. Weinert C. A social support measure: PRQ85. *Nurs Res* 1987;**36**(5):273-7.

- 1
2
3 31. Weinert C, Cudney S, Hill WG. Rural women, technology, and self-management of chronic
4 illness. *Can J Nurs Res* 2008;**40**(3):114-34.
5
6
7
8 32. Long KA, Weinert C. Descriptions and perceptions of health among rural and urban adults
9 with multiple sclerosis. *Res Nurs Health* 1992;**15**(5):335-42.
10
11
12 33. Canada S. Canadian Community Health Survey: Data Liberation Initiative 2010.
13
14 34. Gagne L, Tedds, L., Sullivan, J., Berniaz, K. An introduction to the UT/DLS: Microdata
15 analysis subsetting. Secondary An introduction to the UT/DLS: Microdata analysis
16 subsetting n.d. <http://sda.chass.utoronto.ca/sdaweb/doc/SDAtutorial2008.pdf>.
17
18
19 35. Hawthorne G, Elliott P. Imputing cross-sectional missing data: comparison of common
20 techniques. *Aust N Z J Psychiatry* 2005;**39**(7):583-90.
21
22
23 36. Evans C, Beland SG, Kulaga S, et al. Incidence and prevalence of multiple sclerosis in the
24 Americas: a systematic review. *Neuroepidemiology* 2013;**40**(3):195-210.
25
26
27 37. Canada S. Population by sex and age group Secondary Population by sex and age group
28 2012. <http://www.statcan.gc.ca/tables-tableaux/sum-som/l01/cst01/demo10a-eng.htm>.
29
30
31 38. Buchanan RJ, Chakravorty BJ, Tyry T, et al. Age-related comparisons of people with
32 multiple sclerosis: demographic, disease, and treatment characteristics.
33 *NeuroRehabilitation* 2009;**25**(4):271-8.
34
35
36 39. Giordano A, Ferrari G, Radice D, et al. Self-assessed health status changes in a community
37 cohort of people with multiple sclerosis: 11 years of follow-up. *Eur J Neurol*
38 2013;**20**(4):681-8.
39
40
41 40. Asano M, Duquette P, Andersen R, et al. Exercise barriers and preferences among women
42 and men with multiple sclerosis. *Disabil Rehabil* 2013;**35**(5):353-61.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

41. Kasser SL, Kosma M. Health beliefs and physical activity behavior in adults with multiple sclerosis. *Disabil Health J* 2012;**5**(4):261-8.

42. Jensen MP, Molton IR, Gertz KJ, et al. Physical activity and depression in middle and older-aged adults with multiple sclerosis. *Disabil Health J* 2012;**5**(4):269-76.

43. Krokavcova M, Nagyova I, van Dijk JP, et al. Mastery, functional disability and perceived health status in patients with multiple sclerosis. *Eur J Neurol* 2008;**15**(11):1237-44.

44. Kayes NM, McPherson KM, Schluter P, et al. Exploring the facilitators and barriers to engagement in physical activity for people with multiple sclerosis. *Disabil Rehabil* 2011;**33**(12):1043-53.

45. Chretien JP, Chu LK, Smith TC, et al. Demographic and occupational predictors of early response to a mailed invitation to enroll in a longitudinal health study. *BMC Med Res Methodol* 2007;**7**:6.

46. Manouchehrinia A, Tench CR, Maxted J, et al. Tobacco smoking and disability progression in multiple sclerosis: United Kingdom cohort study. *Brain* 2013;**136**(Pt 7):2298-304.

47. D'Hooghe M B, Nagels G, De Keyser J, et al. Self-reported health promotion and disability progression in multiple sclerosis. *J Neurol Sci* 2013;**325**(1-2):120-6.

48. Marrie RA, Horwitz RI, Cutter G, et al. Smokers with multiple sclerosis are more likely to report comorbid autoimmune diseases. *Neuroepidemiology* 2011;**36**(2):85-90.

49. Weiland TJ, Hadgkiss EJ, Jelinek GA, et al. The association of alcohol consumption and smoking with quality of life, disability and disease activity in an international sample of people with multiple sclerosis. *J Neurol Sci* 2014;**336**(1-2):211-9.




Figure 1: Survey Respondents
161x152mm (300 x 300 DPI)



Figure 2: Number of respondents mapped by postal code
26x15mm (300 x 300 DPI)




Figure 3: Frequency of Barthel Index score categories: A score of 100 on the Barthel Index suggests that an individual is independently ambulatory, can ascend and descend stairs on their own, is continent and able to perform ADLs without help. Fifteen percent of this cohort were independent. A Barthel Index score of 91-99 suggests slight dependence (16%); 61-90 suggests moderate dependence (47%); 21-60 suggests severe dependence (16%) and 0-20 represents complete dependence (7%).
161x152mm (300 x 300 DPI)




Figure 4: Frequency of Frenchay Activities Index score categories
174x139mm (300 x 300 DPI)

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Complete
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	✓
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	✓
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	✓
Objectives	3	State specific objectives, including any prespecified hypotheses	✓
Methods			
Study design	4	Present key elements of study design early in the paper	✓
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	✓
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up <i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	✓
		(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed <i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	✓
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	✓
Bias	9	Describe any efforts to address potential sources of bias	✓
Study size	10	Explain how the study size was arrived at	✓
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	✓
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	✓
		(b) Describe any methods used to examine subgroups and interactions	✓
		(c) Explain how missing data were addressed	✓
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	✓
		(e) Describe any sensitivity analyses	✓

Continued on next page

Results

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	✓
		(b) Give reasons for non-participation at each stage	✓
		(c) Consider use of a flow diagram	
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	✓
		(b) Indicate number of participants with missing data for each variable of interest	✓
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	✓
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	✓
		(b) Report category boundaries when continuous variables were categorized	✓
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	✓
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	✓

Discussion

Key results	18	Summarise key results with reference to study objectives	✓
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	✓
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	✓
Generalisability	21	Discuss the generalisability (external validity) of the study results	✓

Other information

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	✓
---------	----	---	---

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.